

Three Dollars per Annum, in Advance. Single Copies, Twenty-Five Cents.

Prof. Hadd

VOL. III.

No. V.

THE

MATHEMATICAL MONTHLY.

FEBRUARY, 1861.

EDITED BY

J. D. RUNKLE, A.M., A.A.S.

CAMBRIDGE:

PUBLISHED BY SEVER AND FRANCIS.

NEW YORK:

A. S. BARNES AND BURR.

LONDON: TRÜBNER & CO.

1861.

Agassiz's Natural History,

VOL. III.

LITTLE, BROWN, & CO.,

110 Washington Street, Boston,

HAVE JUST PUBLISHED

Contributions to the Natural History

OF THE

UNITED STATES OF AMERICA.

BY LOUIS AGASSIZ.

Vol. III. 4to. 26 Plates. \$12.00.

This volume is divided into three parts : Part I. Acalephs in General; Part II. Ctenophore; Part III. Discophore. It contains a history of our knowledge of these animals from the earliest period to the present time; their classification by different naturalists; with a full account of their structure and mode of development, from the egg, through all their metamorphoses, to their full growth; with remarks upon their habits and appearance along our coast. It is very fully illustrated by twenty-six beautifully executed plates, each of which is crowded with figures.

In this volume the views of Darwin on the Origin of Species are also considered.

To Advertisers.

IOWA SCHOOL JOURNAL.

EDITED BY

HON. THOS. H. BENTON, JR.,

Sec'y of the State Board of Education.

The second Volume of the IOWA SCHOOL JOURNAL offers, by reason of the new arrangements under which it will be conducted, and the largely increased number of subscribers that will not fail to be the result, the best medium for the class of advertising that seeks Educational Papers, to be found in the West.

The connection of the Secretary of the Board of Education, ex-officio Superintendent of Public Instruction, with the Journal, will give it advantages possessed by no other similar publication in the country.

We shall be pleased to receive the favors of advertisers at low rates.

ADDRESS

MILLS BROTHERS,

Publishers Iowa School Journal,

Des Moines, Iowa.

Contents.

FEBRUARY, 1861.

	PAGE		PAGE
PRIZE PROBLEMS FOR STUDENTS	129	SQUARES OF NUMBERS ENDING IN 5, 25, AND 75.	
REPORT OF THE JUDGES UPON THE SOLUTIONS OF		By J. F. Roberson, State Univ., Ind.	150
THE PRIZE PROBLEMS IN NO. I., VOL. III.	130	SOLUTION OF PROBLEM V., No. X., VOL. II. By	
DESCRIPTIVE GEOMETRY OF ONE PLANE. By Wil-		Asher B. Evans, Madison Univ.	151
liam Watson.	133	ON AN APPROXIMATE AND GRAPHICAL RECTIFICA-	
NOTES AND QUERIES	149	TION OF THE CIRCLE. By A. S. Herschel.	152
A GENERAL METHOD OF FINDING TANGENTS TO AL-		MATHEMATICAL MONTHLY NOTICES	155
GEBRAIC CURVES. By Walter Holliday, Univ. of Va.	149	EDITORIAL ITEMS	160

Terms to Subscribers and Advertisers.

No. of Copies to one Address.	Volume.	Terms.	Premium to Clubs.	Advertisement Pages.	First Insertion.	Subsequent Insertions.
1	III.	\$ 3.00		1	\$ 15.00	\$ 12.00
2	III.	5.00		$\frac{1}{2}$	9.00	7.00
3	I., II., III.	6.00		$\frac{1}{3}$	5.00	4.00
5	III.	11.00	III.	$\frac{1}{5}$	3.00	2.00
10	III.	20.00	II., III.			
15	III.	30.00	I., II., III.			

Parties in New York wishing to advertise in the Mathematical Monthly will please send their favors to Messrs. A. S. BARNES AND BURR by the 15th of each month. All club-subscriptions and business communications must be addressed to

SEVER AND FRANCIS, Publishers,

Cambridge, Mass.

University Press, Cambridge : Electrotyped and Printed by Welch, Bigelow, & Co.



THE

MATHEMATICAL MONTHLY.

Vol. III. . . . FEBRUARY, 1861. . . . No. V.

PRIZE PROBLEMS FOR STUDENTS.

I. FIND a point without two concentric circles, from which if two tangents be drawn to the circles the one shall be the double of the other.

II. Any whole number and its fifth power, when divided by 30, leave the same remainder; also, the number and its seventh power, when divided by 42, leave the same remainder. — Communicated by ASHER B. EVANS, Madison University, Hamilton, N. Y.

III. There are n straight lines making, with another fixed straight line, the angles $\alpha, \beta, \gamma \dots$; a point, P , is taken, such that the sum of the squares of the perpendiculars from it on these n lines is constant. Find the conditions that the locus of P may be a circle. — Communicated by GEORGE B. HICKS, Cleveland, Ohio.

IV. If A and B represent the semi-axes of an ellipse, the altitudes of the minimum circumscribing isosceles triangles, having their vertices in the axes produced, are as $A : B$. — Communicated by JAMES F. ROBERSON, Grantsburgh, Indiana.

V. Prove that, of all circular sectors having the same perimeter, the one of greatest area is that in which the circular arc is double the radius.

Solutions of these problems must be received by April 1st, 1861.

REPORT OF THE JUDGES UPON THE SOLUTIONS OF THE
PRIZE PROBLEMS IN No. I., Vol. III.

THE first Prize is awarded to A. G. BARKER, Waterville College, Me.

The second prize is awarded to H. TEJMAN, Baltimore, Md.

The third prize is awarded to E. O. GIBSON, South New Berlin, N. Y.

PRIZE SOLUTION OF PROBLEM I.

By E. K. LEONARD, Marietta College, Ohio.

Let AB and CD be two diameters of a given circle, drawn at right angles to each other; AEB a circular arc described with radius DA or DB . Prove that the area of the lune $AEB C$ equals the area of the triangle ADB .

Let R be the radius of the arc AEB , and r that of the circle. When $ABC = \frac{1}{2} \pi r^2$, $ADB C = \frac{1}{4} \pi R^2$, $r : R = 1 : \sqrt{2}$. Substituting 1 and $\sqrt{2}$ for r and R , we have $ABC = ADB E$. Subtracting ABE from each, we get $AEB C = ADB$.

Most of the solutions of this problem are nearly the same as the above.

PRIZE SOLUTION OF PROBLEM II.

By E. O. GIBSON, South New Berlin, N. Y.

Of all isoperimetrical polygons having the same number of sides, the greatest is that which is equilateral.

Since an isosceles triangle has a greater area than any other triangle of equal base and perimeter, it follows that if any polygon, $ABCD$, &c., be a maximum among others of equal perimeters and the same number of sides, all the triangles, ABC , BCD , &c., must be isosceles, and have $AB = BC = CD = \&c.$

PRIZE SOLUTION OF PROBLEM III.

In any triangle, if P , P' , and P'' denote the perpendiculars from the vertices to the opposite sides; p , p' , and p'' the portions of the same between their point of intersection and the sides. Prove that

$$\frac{p}{P} + \frac{p'}{P'} + \frac{p''}{P''} = 1.$$

Mr. G. A. OSBORNE, Jr., says: Let a, b, c denote the sides of the triangle perpendicular respectively to P, P', P'' . Then

$$ap + bp' + cp'' = 2 \text{ (area of triangle),}$$

and $aP = bP' = cP'' = 2 \text{ (area of triangle).}$

Dividing each term of the first equation by the term below it, gives

$$\frac{p}{P} + \frac{p'}{P'} + \frac{p''}{P''} = 1.$$

Mr. GIBSON notices that, in an obtuse-angled triangle, as the parts of the perpendiculars let fall on the sides produced, and which are represented by p and p' , lie beyond those sides, they may be regarded as minus, and we still have

$$\frac{p}{P} + \frac{p'}{P'} + \frac{p''}{P''} = 1.$$

Mr. G. B. HICKS remarks, that this proposition may be included in the still more general statement, that, if from the angles of a triangle three lines, Aa, Bb, Cc , be drawn to meet the opposite sides, and from any point, O , within the triangle three lines, Oa, Ob, Oc , be drawn parallel to them to meet the sides, then

$$\frac{Oa}{Aa} + \frac{Ob}{Bb} + \frac{Oc}{Cc} = 1.$$

PRIZE SOLUTION OF PROBLEM IV.

By A. G. BARKER, Waterville College, Maine.

At what angle must the rudder of a vessel be inclined to the stream, that the effect produced may be a maximum?

Let the angle made by the rudder with the direction of the vessel be denoted by φ , the horizontal distance of the vessel's centre of gravity from the line of contact of the rudder by a , and the length of the rudder by b . The force exerted by the water upon the rudder parallel to the direction of the vessel's motion is proportional to $\sin \varphi$, since this force is zero for φ equal zero, and a maximum for φ

equal 90° . We may, therefore, take $\sin \varphi$ as its measure ; and since φ is the angle this force makes with the rudder, its component perpendicular to the rudder is $\sin^2 \varphi$, which we may suppose to act in the perpendicular passing through its middle point. This perpendicular produced will meet the line of direction of the vessel at a distance $\frac{\frac{1}{2}b}{\cos \varphi}$ from the line in which the rudder joins the vessel, and $a + \frac{\frac{1}{2}b}{\cos \varphi}$ is the length of the arm upon which the force $\sin^2 \varphi$ acts at an angle $\frac{1}{2} \pi - \varphi$. The component of $\sin^2 \varphi$ perpendicular to $a + \frac{\frac{1}{2}b}{\cos \varphi}$ is $\sin^2 \varphi \cos \varphi$, and

$$u = \left(a + \frac{\frac{1}{2}b}{\cos \varphi} \right) \sin^2 \varphi \cos \varphi = \max.$$

$$\begin{aligned} \therefore \frac{du}{d\varphi} &= 2a \sin \varphi \cos^2 \varphi - a \sin^3 \varphi + b \sin \varphi \cos \varphi = 0 \\ &= 3a \cos^2 \varphi + b \cos \varphi - a = 0; \end{aligned}$$

$$\therefore \cos \varphi = - \frac{b \pm \sqrt{(12a^2 + b^2)}}{6a}.$$

The second derivative gives

$$\frac{d^2u}{d\varphi^2} = -6a \cos \varphi \sin \varphi - b \sin \varphi,$$

which, for $\varphi < 180^\circ$, is negative for the positive value of $\cos \varphi$; and therefore this value of $\cos \varphi$ will make u a maximum.

The negative value of $\cos \varphi$ gives

$$\frac{d^2u}{d\varphi^2} = \sqrt{(12a^2 + b^2)} \sin \varphi,$$

a positive quantity; and u is therefore a minimum for the negative value of $\cos \varphi$.

When $b = 0$, that is, when the length of the rudder is neglected, $\cos \varphi = \pm \sqrt{\frac{1}{3}}$; $\therefore \varphi = 54^\circ 44'$, or $125^\circ 16'$.

None of the solutions of Problem V. are thought worthy of publication.

SIMON NEWCOMB,
W. P. G. BARTLETT,
TRUMAN HENRY SAFFORD.

DESCRIPTIVE GEOMETRY OF ONE PLANE.

CHAPTER I.

THE POINT, THE RIGHT LINE, AND THE PLANE.

THE POINT.

§ 1. A POINT in space is represented by its orthographic projection upon a fixed plane, called the *plane of reference*, and a number written near the projection, to show the distance of the point from this plane: this number is called the *reference* of the point with respect to the plane.

The plane of reference will at present be considered horizontal, and entirely below the magnitudes to be represented, thus avoiding any confusion which might arise from the use of negative references.

THE RIGHT LINE.

§ 2. A right line in space is determined by the projections and corresponding references of two of its points.

§ 3. PROBLEM I. *Given the projection of a point situated upon a given right line, to find the reference of the point.*

Let AB (Fig. 2) be the projection of a line, and let the references of two of its points, projected at A and B , be respectively α and β . It is required to find the reference of the point, of which C is the projection. If we imagine the line in space revolved about its horizontal projection as an axis, so as to coincide with the plane of reference, it will take the position $A'B'$, making AA' equal to α , and BB' equal to β . Calling the unknown reference of C , γ , we have

$$BA : BC = \alpha - \beta : \gamma - \beta,$$

whence

$$\gamma = \frac{BC}{BA} (\alpha - \beta) + \beta.$$

The lengths of AB and BC are known, having been measured in units of the scale (Fig. 1), and α and β are measured upon the same scale.

Conversely, if it be required to find the projection of that point of the line whose reference is γ , we have, from the same proportion,

$$BC = AB \left(\frac{\gamma - \beta}{\alpha - \beta} \right).$$

Or the point C' may be found by laying off on the line AA' a distance AD equal to γ , and drawing DC' parallel to AB .

§ 4. PROB. II. *To find the distance between two given points.*

Let A and B (Fig. 2) be the given points, then evidently the distance required will be $\sqrt{AB^2 + (\alpha - \beta)^2}$.

§ 5. *Definitions.* — The *tangent* of the angle made by the line with the plane of reference, $\left(\frac{\alpha - \beta}{AB} \right)$, is called the *declivity of the line*, and when put in the form $\frac{1}{m}$, it shows that points of the line whose projections are m units apart differ in reference by unity.

The distance between the projections of two points is called their *interval*; and the distance between the projections of two points of a line differing from each other by a unit of reference is called the *interval of the line*.

§ 6. By an application of the process explained in Problem I, it is easy to divide the projection of any given line into intervals; the projection thus divided is called the *scale of declivity of the given line*.

§ 7. PROB. III. *To construct the scale of declivity of a given line.*

Let the line be given by its projection, and the references 10.83 of the point A and 8.28 of B (Fig. 3); the interval of the points A and B , measured upon the scale (Fig. 1) is 5.75.

Therefore the declivity of the line is

$$\frac{10.83 - 8.28}{5.75} = \frac{2.55}{5.75} = \frac{1}{2.25} = \frac{1}{m}.$$

Therefore the interval of the line is $m = 2.25$. Therefore, setting off a distance 2.25 from A towards B , we have the point C , of which the reference is less by unity than that of A , i. e., 9.83.

But it is usual to divide the projection into integer references. To find the point 10, set off from A towards B a distance $2.25 \times 0.83 = 1.87$ of the scale (Fig. 1); from 10 set off a distance 2.25, which gives the point 9, &c. Divide one of the intervals, as 9–8, into 10 equal parts, and the scale will be completed.

§ 8. PROB. IV. *Through a given point to draw a line parallel to a given line.*

Let the line ab be given by the projections and references of two points, 12 of a and 7.80 of b , and the point by its projection c and its reference 8.50. Draw through c a line parallel to ab , and lay off on it from c the interval ab ; this gives a point d of the required line, of which the reference is $8.50 + 4.20 = 12.7$.

THE PLANE.

§ 9. The planes which are required to be represented are of two kinds, those which are limited in extent, as faces of polyhedrons, and those which are unlimited.

An unlimited plane is determined by the position of one of its lines of greatest declivity, that is, one of the lines of the plane making the maximum angle with the horizontal plane of reference; this line is evidently perpendicular to every horizontal line of the plane so defined.

The scale of declivity of this line is called the *scale of declivity of the plane*, and to distinguish it from that of an ordinary line it is made double.

A limited plane is represented by the projection of its perimeter, to which is added either its scale of declivity or the references of its angular points. Often a plane is given by two points of its line of greatest declivity.

§ 10. PROB. V. *To pass a plane through three given points.*

Let a , b , c (Fig. 5) be the points, of which the references are respectively 10.5, 7, and 4. Join a and c , and find on ac the point d , whose reference is the same as b (7); join d and b . This will be a horizontal line of the required plane; the line of maximum declivity will be perpendicular to it.

To find another point, draw through c a line parallel to db ; it will be the horizontal (4), which determines the line of greatest declivity.

If we have the projection of any point, as e , situated upon this plane, to find its reference, it will be sufficient to drop a perpendicular from e to the scale, and find by Prob. I. the reference of the point of intersection, which will be the reference required.

§ 11. PROB. VI. *To find the intersection of two planes.*

Let the two planes M and N (Fig. 6) be given by their scales of declivity.

In one plane, M , draw any two horizontals; through 7 and 11, for instance. In the other plane, N , draw horizontals having the same references, and the points a and b , their intersections, are two points of the line of intersection of the two planes.

It may happen that the horizontals of each plane are nearly or quite parallel. In the first case, take two auxiliary planes, Q and P (Fig. 7). Q intersects M in the line xy , and also N in the line vt ; therefore a , the intersection of xy and tv , is a point common to the three planes M , N , and Q , or a point of the line of intersection of M and N . The same construction with P gives the point b ; whence ab is the line of intersection required.

If the horizontals were exactly parallel, one auxiliary plane would be sufficient, since the line of intersection would be a horizontal.

§ 12. PROB. VII. *Through a given point in a given plane, to draw a line of the plane having a given declivity.*

Let M (Fig. 8) be the given plane, a the projection of the given

point, and 2.7 its reference; let the declivity of the required line be $\frac{6}{10}$. Draw any horizontal of M , as 6, and take the difference between its reference and that of the given point ($6 - 2.7 = 3.3$); multiply this difference by the reciprocal of the declivity of the required line ($3.3 \times \frac{10}{6} = \frac{11}{2} = 5.5$); then with the projection of the given point as a centre, and a radius 5.5, describe an arc cd ; and where it meets the horizontal 6 will be another point of the required line.

There will be two solutions, if the declivity of the line be less than that of the plane; one, if it be equal, none, if it be greater.

§ 13. PROB. VIII. *Through a given line pass a plane of a given declivity.*

Let the declivity of the required plane be $\frac{1}{m}$ (Fig. 9). Conceive any point of the line, as b , to be the vertex of a cone, of which the elements have a declivity $\frac{1}{m}$; to trace the base of the cone, it will be sufficient to take a radius, m , and with b as a centre describe a circle; the plane of this base will be horizontal, whose reference is less by unity than that of b . Now, from the point c upon the given line, whose reference is less than b by unity, draw a tangent to the circle; this tangent will be a horizontal of the required plane. Draw a parallel through b , and we have another, whence its line of declivity is immediately determined. In general there will be two solutions.

Sometimes it happens, in the solution of this problem, that the line being very little inclined, the position of the second point would lie off the paper. The solution for this case is to take any other point, d , of the line, and consider it also the vertex of a cone having its base on the same horizontal plane, and its declivity the same, the radius of its base would be m times the altitude; a horizontal of the plane would then be found by drawing a tangent common to the two circles of the base. (Fig. 10.)

If the line given were exactly horizontal, then one cone would be sufficient.

§ 14. PROB. IX. *To find the point in which a line pierces a plane.*

Let M (Fig. 11) be the given plane, and A the given line. Through the line A pass any arbitrary plane, X , and find its intersection, mn , with M ; the point in which this line of intersection meets A is the point sought.

This method may be used to ascertain the point in which two lines having the same projection intersect.

Pass through the two given lines any two arbitrary planes, X and Y , and determine their intersection; the point in which this intersection meets the given lines is their common point. Or the point of intersection may be found by revolving the lines about their common projection as an axis until they are parallel to the horizontal plane of reference. [The projection may be considered as situated in any horizontal plane, since, as in this case, it is often convenient to work upon horizontal planes of considerable references.]

Fig. 12 represents these two solutions.

§ 15. PROB. X. *From a given point in space to drop a perpendicular upon a given plane, to find the point in which it pierces the plane and the length of the perpendicular.*

First Solution. — The projection of the perpendicular will evidently be parallel to the scale of declivity of the plane, and its declivity the reciprocal of that of the plane.

Suppose the declivity of the plane is $\frac{1}{m}$, then the interval upon the perpendicular will be $\frac{1}{m}$.

Therefore, through the projection of the given point a draw a line parallel to the scale of declivity of the plane, and lay off from a on this line intervals of $\frac{1}{m}$, measured upon the scale of equal parts (Fig. 1); the references of the perpendicular will increase in a direction contrary to that of the scale of declivity of the plane.

The point in which the perpendicular pierces the given plane may be found by § 16, and its length by § 4.

If the line of declivity of the plane be given by the references of two of its points, c and d , the perpendicular can still be drawn without constructing the scale of the plane. For after drawing the parallel to the line of declivity of the plane, set off on it from the point a a distance equal to the difference of the references of c and d , and the reference of this last point will be the reference of a *plus* or *minus* the length of the interval between c and d , according as it is laid off *above* or *below* the point a . That is, intervals of the plane correspond to differences of reference on the perpendicular, and reciprocally.

There is another solution to this problem, which may be used to advantage in many of the subsequent problems.

Let a (Fig. 13) be the given point, de a section of the plane of reference, dc the line of maximum declivity of the given plane, ac the required perpendicular, and de its horizontal projection; $\tan cde = \frac{1}{m}$, the declivity of the given plane. Then the length of

$$ac = ad \cos \tan^{-1} \frac{1}{m}, \quad \text{and} \quad de = ad \sin \tan^{-1} \frac{1}{m} \cos \tan^{-1} \frac{1}{m};$$

these two may be readily constructed in connection with the scale of declivity of the plane.

Let M (Fig. 14) be the given plane, and a be the projection of the point of which the reference is 11. Let fall from a a perpendicular to the scale, and find the corresponding reference 8.4 of b . From b set off a distance bc equal to an interval of the scale M , and also set off bd equal to a unit of the scale of equal parts (Fig. 1); join c and d , and produce this line. From c set off a distance ce equal to 2.6, the difference of reference between b and a . Draw ef perpendicular to cd , and fh parallel to ce ; and the point h , where the last line cuts the parallel through a to the scale M , is the point sought. ah is the

projection of the required perpendicular and cf is its length; the angle fce is the angle of declivity of the plane.

The interval ah may be easily calculated. Let the declivity of the plane be $\frac{1}{m}$, and the difference of reference between the points a and b be denoted by δ ; then the length of the required perpendicular is

$$p = \delta \cos \tan^{-1} \frac{1}{m} = \delta \cdot \frac{m}{\sqrt{(m^2+1)}},$$

$$ah = \delta \cdot \cos \tan^{-1} \frac{1}{m} \sin \tan^{-1} \frac{1}{m} = \delta \frac{m}{\sqrt{(m^2+1)}} \cdot \frac{1}{\sqrt{(m^2+1)}} = \delta \frac{m}{m^2+1}.$$

The difference of reference between the point h and the horizontal ab is $\delta \sin^2 \tan^{-1} \frac{1}{m} = \delta \frac{1}{1+m^2}$.

§ 16. PROB. XI. *To draw through a given point a plane perpendicular to a given line.*

The projection of the scale of declivity of the plane must be parallel to that of the given line, and its declivity is the reciprocal of that of the line.

So the scale is found in the same manner as that of the required line by the first method of the last problem.

§ 17. PROB. XII. *To find the shortest distance from a point, a , in space to a line, A .*

First Method. — Pass through a a plane perpendicular to A , and find the point in which A pierces this plane; the line connecting this last point with a will be the perpendicular required.

Second Method. — Let a (Fig. 15) be the given point, and A the given line; join A and the point b of A having the same reference (7), ab will be a horizontal of their plane (X); rotate this plane about ab as an axis until it becomes parallel to the horizontal plane of reference; any point of A , as c , will be found at c' , by taking sc' equal to the distance between the axis and c in space. Therefore A' will be the revolved position of A ; ad' will be the length of the

perpendicular, and, making the counter revolution, we have $a d$, its projection.

§ 18. PROB. XIII. *Through any two lines, A and B, pass two parallel planes.*

Through any point of A draw a line parallel to B ; these two lines will determine the first plane. Then through any point of B draw a plane parallel to the first.*

§ 19. PROB. XIV. *Through a given line, A, to pass a plane perpendicular to a given plane, M.*

From any point of A drop a perpendicular upon M ; this perpendicular and the line A determine the required plane.

§ 20. PROB. XV. *To measure the angle of two lines, A and B.*

Draw through any point of A (Fig. 17) a line parallel to B , as ay , and pass a plane through A and ay . Rotate this plane about any horizontal xy , so as to be parallel to the plane of reference. The new positions of the lines determine the angle sought, $x a' y$.

Since the three sides of the triangle xay are very easily determined, the angle may be found without rotation by the formula,

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}.$$

§ 21. PROB. XVI. *Through a given point, a, to draw a line which shall make a given angle, μ , with a given line, A.*

Join a (Fig. 18) and the point of A having the same reference, this will be a horizontal of their common plane; rotate this plane till it is parallel to the plane of reference, and find A' , the new position of the line A . Draw through a a line $a c'$, making the given angle μ with the line A' , and make the counter revolution: ac will be the line sought.

If the tangent of the angle μ be given, $\frac{1}{m}$, the point c' may be

* The construction for drawing one plane parallel to another is the same as for drawing one line parallel to another, and was explained by Prob. IV. Fig. 16 gives the construction.

found by dropping from a the perpendicular ad' , and measuring off the distance $d'c'$ equal to m times ad' .

§ 22. PROB. XVII. *To construct the angle of a given line, A , and a plane, M .*

From any point of A drop a perpendicular upon M , and by Prob. XV. find the angle between A and this perpendicular; this will be the complement of the angle sought.

§ 23. PROB. XVIII. *To find the scale of declivity of a line, knowing its projection, and also knowing it to be perpendicular to a given line, A , at a given point, a .*

Through the angular point a draw a plane, X , perpendicular to the given line. The unknown line will be in this plane; therefore drawing in X any horizontal, the point in which it meets the given projection of the unknown line will correspond to a point of the required line of the same reference; whence its scale may be immediately constructed.

§ 24. PROB. XIX. *To find the scale of declivity of a line, knowing its projection, and also the angle, μ , which it forms with a given line at a given point.*

If with the given angular point as a centre a sphere be conceived to be described, the four points of its surface in which the two lines and their two projections pierce it will form the angular points of a spherical quadrilateral, a, b, c, d (Fig. 19), in which ab is the angle of declivity of the given line, ad the given angle μ , cb that of the projections, abc, bcd are right angles, and dc is the unknown angle of declivity: whence the side dc may be calculated by the rules of spherical trigonometry; but the graphic solution is as follows:—

Let a (Fig. 20) be the angular point, ab the projection of the given line, ae that of the unknown line. Pass a plane through the given line and the projection of the unknown line; its scale will be M , its angle of declivity Mfg , the complement of which is gfe . Rotate

the line ab about ae as an axis, so as to coincide with the horizontal plane through a ; it will take the position ac .

Make the angle cad equal to the given angle μ . Now, comparing Figs. 19 and 20, we have the three parts of the spherical triangle acd (Fig. 19), ac , acd , and ad , which correspond respectively to eac , gfe , and cad of (Fig. 20); and it is required to find the side dc (Fig. 19), or, in other words, the other face of the pyramid. To do this, erect at any point m of ae (Fig. 20) a perpendicular, mn ; make the angle mnr equal to gfe ; then mr is a line of the unknown face revolved about nm as an axis, to coincide with the horizontal plane. At n erect a perpendicular to mn , and also draw nv perpendicular to ac , and ns perpendicular to nv and equal to nr . The line nvx is the horizontal trace of a vertical plane; and joining s and x , sx is the revolved position of the intersection of the unknown face and the vertical plane ux . With v as a centre and uv as radius describe an arc ut , cutting sx in t . Let fall from t the perpendicular tz to ux , and from z a perpendicular, zy , to ae . With x as a centre, and a radius xt , describe an arc meeting zy produced in t' . Join t' and a , and we have eah , the angle of declivity sought, and ah , the position of the line required revolved about its horizontal projection.

§ 25. PROB. XX. *To reduce an angle to the horizon; that is, having given μ the angle which a line forms with a given line at a certain point, and also its declivity, to find its projection.*

Let ab (Fig. 21) be the given line, and a the angular point. Revolve the line about its projection so as to coincide with the horizontal plane through b ; it will take the position ba' . Make the angle $a'a'c$ equal to the angle of declivity of the unknown line, and $ba'd$ equal to μ , the given angle.

$a'e$ intersects ab in c , therefore with a as a centre, and a radius ac , describe a circle; it will be the base of a cone of which a is the vertex, and every element of this cone will have the given declivity.

Take $a'd$ equal to $a'c$, and with b as a centre and bd as a radius describe an arc intersecting the former in the points m and n ; am or an will be the projection required. The line required and the given line are represented in revolved position by $b a' d$, and the third side of the triangle $b a' d$ is horizontal.

§ 26. *Transformation of the Plane of Reference.* — It is convenient often to change the plane of reference and refer the magnitudes to another plane whose scale with respect to the primitive is given; this may be done by an application of Prob. X.

I. *To change with respect to a point, i. e. to find the new projection and reference of a given point, a , with respect to any plane, M .*

Let fall from a a perpendicular upon M ; find the point a' , in which it pierces the plane M , and the length of the perpendicular.

The point a' is the new projection, and the length of the perpendicular is the reference sought.

II. *To change with respect to a line.*

Find the new projections and references of two points of the line. The construction may be simplified by taking as one point that in which the line pierces the plane M , which is its own projection, and has a reference zero.

III. *To change with respect to a plane; change with respect to any three points of the plane.*

The construction is much simplified by finding first the line of intersection of the given plane and the new plane of reference; this line is the new horizontal of zero;* dropping upon this intersection a perpendicular from the new projection of another point of the plane, this will be the projection of the new scale, which is known by the references of two points.

The scale of the primitive with respect to the new plane of refer-

* The term horizontal here is evidently a misnomer, but it cannot lead to any ambiguity.

ence will have the same projection and interval, but will be the negative of it, *i. e.* it will increase from zero in the opposite direction.

The position of the new plane of projection may not be given directly, it may be required to be parallel or perpendicular to a given line or plane. The Problems IV., X., and XI. enable us to find immediately the scale of declivity required.

§ 27. PROB. XXI. *To find the angle of two planes, M and N .*

First Method. — Take the plane M as a new plane of reference; the new angle of declivity of N will be the angle sought. Fig. 22 gives the construction. Find the intersection ce of the two planes M and N ; this is the new horizontal. Let fall from any point of N , as a , a perpendicular upon M ; it pierces it at a' , and its length is bd . Let fall from a' a perpendicular, $a'f$, to ce , the new horizontal; then $a'f$ is the line of declivity of N with respect to M . Erect at a' a perpendicular to $a'f$ equal to bd , the new reference of a' ; join g and f ; fg is the revolved position of the line of maximum declivity of N' , and $a'fg$ is the angle required.

Second Method. — Let fall from any point in space, a , a perpendicular upon each plane, and measure the angle of these perpendiculars by § 20; this will be the supplement of the angle required. Or the length of the perpendicular upon either plane, as M , divided by the distance from its foot to the point in which the perpendicular upon N pierces the plane M , will be the tangent of the angle sought.

Third Method. — Let M and N (Fig. 23) be the two planes, ab their line of intersection. Conceive a plane drawn perpendicular to ab at the point y ; xyz will be a horizontal of this plane, having a reference 70. From y let fall a perpendicular, yy' , upon ab , the revolved position of the line of intersection of the planes, and take yy'' equal to yy' , the angle $xy''z$ will be the angle sought.

§ 28. PROB. XXII. *Through a given right line to pass a plane which shall make a given angle, μ , with a given plane.*

Let A (Fig. 24) be the given line, and M the given plane. Take M as the new plane of reference.

To find the new position, A' , of A , find first the point a , in which A pierces M ; and then from any point of A , as d , let fall a perpendicular upon the plane M ; it pierces this plane at the point c , and its length ex is the new reference of c ; therefore ac is the new projection A' . Denote $\tan \mu$ by $\frac{1}{m}$, the given declivity. Conceive any point of A' , as k , the vertex of a cone, whose elements have the given declivity. The base of this cone is a circle situated in the new plane of reference, having for a centre k , and for a radius the reference of k multiplied by m .

Drawing from a a tangent to this circle, it will be a horizontal of the required plane with respect to the new plane of reference. This new horizontal meets the horizontal of zero of the primitive plane at the point y , therefore y is common to the primitive plane, to the new plane, and to the required plane; it is also its own projection in both planes, and has a reference zero. Therefore, joining y and the point z of the line A , whose reference is zero, we have the line zy , a horizontal of the required plane with respect to the primitive plane. Any other parallel, as through 40, is sufficient to determine the scale of declivity N of the required plane. The problem has in general two solutions. The other plane is determined by the other tangent, ay' , and has for a scale N' .

Second Method.— Let $\frac{1}{m}$ be the tangent of the given angle, A the given line, and M the given plane. From any point of A let fall a perpendicular upon M , and measure its length, p . Measure also the distance from the foot of p to the point in which A pierces M , and call this distance A' .

The distance from the foot of p to the intersection of the unknown plane with m is $m \cdot p$, and the sine of the angle which A' makes with the same intersection is $\frac{m \cdot p}{A'}$.

Through the foot of p pass a plane perpendicular to A' ; and upon its intersection with M , starting from the foot of p , lay off a distance equal to $A' \times \tan \left(\sin^{-1} \frac{p \cdot m}{A'} \right)$, or $\frac{A \cdot p \cdot m}{\sqrt{(A^2 - p^2 \cdot m^2)}}$; and this point will be one of the required planes which, with the given line, immediately determines it.

§ 29. PROB. XXIII. *To determine the perpendicular between two right lines not in the same plane.*

First Solution. — Let A and B (Fig. 25) be the two lines. Pass a plane, X , perpendicular to A at any point, c , and take it as the new plane of reference. The new projection of B is B' , and that of A is reduced to the point c . Draw cr perpendicular to B' ; this will be the perpendicular required. Since cr in space is parallel to the plane X , the new reference of c (the point in which cr meets the new projection of A) is the same as that of r . Therefore, returning to the primitive plane of reference, the point r is projected in t , and making cs in space equal to the new reference of r , st will be the projection of the perpendicular between the lines A and B , and cr will be its length.

Second Direct Solution. — Let A and B (Fig. 26) be the given lines. Pass through B a plane, X , parallel to A . Then from any point of A , as a , let fall a perpendicular to X ; it pierces the plane X in a' , and cd is its length. A line through a parallel to A will be the projection of A upon the plane X . The point n , in which this projection meets B , is one point of the perpendicular sought; drawing mn parallel to aa' , we have the required projection of the perpendicular distance between A and B .

§ 30. PROB. XXIV. *Given a plane, M , and the projection of a limited line, A , of this plane to construct upon A a cube which shall rest upon M .*

Let ad (Fig. 27) be the length, A , of the given projection. Make the angle sxy equal to the angle of declivity of M . Rotate the plane M about any horizontal x , as 50. To find the new position of

d , let fall from d a perpendicular to sx , and produce it to n on the line xy ; then xn will be the distance of d from the axis of rotation, and will make known the side ad' of the cube. The square $ad'e'b'$ will be the base in revolved position, making the counter rotation, the base will be projected in $abcd$. To find the upper base after revolution, make xy equal ad' , and let fall the perpendicular ys ; this will be the length of the horizontal projections of the edges of the cube projected at $abcd$; whence, drawing through a, b, c, d lines perpendicular to the axis of rotation and equal to ys , we have the upper vertices; and the difference of reference between any two points symmetrically situated, as b and h , is sx , whence the cube is completely determined.

§ 31. PROB. XXV. *To find the intersection of two polyhedrons.*

Solution. — Pass a number of horizontal planes through each solid. Then the intersections of horizontals of the same reference, drawn in the intersecting adjacent faces of the two polyhedrons, make known the lines of intersection of those faces.

Fig. 28 represents an application of this method to the determinations of the intersections of two quadrangular pyramids.

§ 32. PROB. XXVI. *To find the shadow of any polyhedron.*

Let $abcde$ (Fig 29) be the lower base, and $a'b'c'd'e'$ the upper base of a polyhedron. Let the reference of the upper base be 80, and that of the lower base 0. Instead of supposing, as usual, the rays of light parallel to the diagonal of a cube resting upon the horizontal plane, let them be supposed to make an angle of 45° with the plane of reference, and have a direction indicated by the arrow A .

To find the shadow of any point, as f' , draw through f' a parallel to A , and lay off a distance $f'f''$ equal to the reference 80 of f' . The same constructions for d' and e' give the points d'' and e'' . Joining d and d'' , f and f'' , we have $ff''e''d''d$ as the shadow required.

NOTES AND QUERIES.

I. *A General Method of finding Tangents to Algebraic Curves.*— If the curve be referred to rectangular axes, and $x_1 y_1$ be the co-ordinates of the point of tangency, the equation of a line passing through this point will be of the form,

$$y - y_1 = a(x - x_1).$$

It is required to find the value of a when this line is tangent to the curve. Refer the curve to a system of polar co-ordinates, whose pole is at the point $x_1 y_1$, and axis parallel to the axis of x . Since the pole is upon the curve, the absolute terms will disappear from this equation. Every term of the equation will now contain r , the first power of which may be divided out. Then making $r = 0$, find the corresponding value of $\tan \theta$, which will evidently be the required value of a . This value of $\tan \theta$ may always be found by the solution of an equation of the first degree. For any term that contains the product of $\sin \theta$ and $\cos \theta$, or the higher powers of either, will also contain the higher powers of r , and will therefore disappear when $r = 0$. Hence it readily appears that in practice we may neglect all terms except those which contain the first power of r ; and from these we shall find the required value of $\tan \theta$.

This method will readily suggest its own demonstration.

EXAMPLES.

1. Take the folio of Descartes, of which the equation is

$$x^3 - 3cxy + y^3 = 0.$$

The equations of transformation are

$$x = x_1 + r \cos \theta, \qquad y = y_1 + r \sin \theta.$$

Hence we have

For numbers ending in 25 and 75 let h represent the hundreds, then 25 and 75 will be represented by $\frac{1}{4}$ and $\frac{3}{4}$; annexing each of these fractions to h , we have the two binomials $(h + \frac{1}{4})$ and $(h + \frac{3}{4})$; squaring each and factoring, they become

$$[h(h + 5) + \frac{1}{16}], \quad [(h + 1)(h + 5) + \frac{1}{16}],$$

each of which may be readily expressed in the form of a rule, by observing that the square of numbers ending in 25 or 75 always ends in 625. Thus,

$$(225)^2 = 50625, (725)^2 = 525625, (375)^2 = 140625, (975)^2 = 950625.$$

Let $h = 100$ and $t = 10$, then we have, for the powers of the above numbers,

$$\begin{aligned} (225)^2 &= (\frac{9}{4}h)^2 = \frac{81}{16}h^2 = 50625, \\ (375)^2 &= (\frac{15}{4}h)^2 = \frac{225}{16}h^2 = 140625, \\ (225)^3 &= (\frac{9}{4}h)^3 = \frac{729}{64}h^3 = \frac{729}{8 \times 8}h^3 = \frac{91125}{8} = 11390625, \\ (25)^4 &= (\frac{5}{2}t)^4 = \frac{625}{16}t^4 = 390625, \\ (15)^6 &= (\frac{3}{2}t)^6 = \frac{729}{64}t^6 = \frac{729}{8 \times 8}t^6 = \frac{91125}{8} = 11390625. \end{aligned}$$

Numbers may also be multiplied together in the same manner. Thus,

$$\begin{aligned} (15 \times 25 \times 5) &= (\frac{3}{2}t \times \frac{1}{4}h \times \frac{1}{2}t) = \frac{3}{16}t^2h = \frac{3}{16}h^2 = 1875, \\ (35 \times 75 \times 65 \times 5) &= (\frac{7}{2}t \times \frac{3}{4}h \times \frac{13}{2}t \times \frac{1}{2}t) = \frac{273}{8 \times 4}t^3h = \frac{34125}{4} = 853125. \end{aligned}$$

— JAMES F. ROBERSON, State University, Bloomington, Indiana.

III. *Solution of Problem V., No. X., Vol. II.* — Suppose the sphere to be divided into an infinite number of concentric spherical shells; let A be their common centre, and B the exterior point; and M any particle on the surface of one of these shells, $r = AM$, $c = AB$, $y = BM$, dr = the thickness of the shell, and ρ = its density.

Since dr is infinitely small, ρ may be considered constant for the

same shell, but varying for *different* shells. The resultant attraction of the whole shell on

$$B = \frac{\pi \rho r dr}{c^2} \int_{c-r}^{c+r} \left(1 + \frac{c^2 - r^2}{y^2}\right) dy = \frac{4\pi \rho r^2 dr}{c^2}.$$

To obtain the part of the shell which exerts half this attraction, take $c - r$ and x for the limits of y , and put

$$(1) \quad \frac{\pi \rho r dr}{c^2} \int_{c-r}^x \left(1 + \frac{c^2 - r^2}{y^2}\right) dy = \frac{2\pi \rho r^2 dr}{c^2}.$$

But $\int_{c-r}^x \left(1 + \frac{c^2 - r^2}{y^2}\right) dy = x - \frac{c^2 - r^2}{x} + 2r$; therefore equation (1) gives $x - \frac{c^2 - r^2}{x} + 2r = 2r$, or

$$(2) \quad x = \sqrt{(c^2 - r^2)}.$$

If C be on the surface of the shell, and $BC = x$, the triangle ABC will be right angled at C . If we suppose r in equation (2) to take successively all values between zero and the radius of the sphere, the triangle ABC will be right angled in every case; hence the locus of the point C , which is always on the bisecting surface, is the surface of the sphere described on AB as diameter. — ASHER B. EVANS, Madison University, Hamilton, N. Y.

IV. *On an Approximate and Graphical Rectification of the Circle.** — The following treatment of this oft-attempted problem is founded upon the singularly close approach that is made by the angle $\tan^{-1}(\frac{1}{4}\pi)$ to a root of the equation $\sec x = \cot x$. The tabular solution of this equation is presented in the form


$$\cos x = \tan x = \sqrt{\left(\frac{\sqrt{(5)} - 1}{2}\right)} = 0.7863,$$

while $\frac{1}{4}\pi = 0.7854$; so that $x = 38^\circ 10' 46''$, and $\tan^{-1} \frac{1}{4}\pi = 38^\circ 8' 46''$:

* From the London Quarterly Journal of Pure and Applied Mathematics, October, 1860.

from which it is clear that if a simple construction can be found for obtaining the angle x that satisfies this equation, the tangent of that angle will yield a very close approximation to the desired periphery of a circle.

Let the chord ADC (Fig. 1) cut the semicircle ADB in D , and the tangent BC in C , in such a manner that $AD = BC$; then shall AD , BC represent *quam proximé* quadrants upon the semicircle ADB . By similar triangles, $\frac{AD}{AM} = \frac{AB}{AD} = \frac{AB}{BC}$, by hypothesis. Or $\sec DAB = \cot DAB$, and DAB is such an angle as x ; therefore



$$BC = AB \tan DAB = 2 \tan x, \text{ if the radius be unity;} \\ = 1.5727; \text{ and } \frac{1}{2} \pi = 1.5708.$$


Therefore BC exceeds the quadrant of ADB by one thousandth part nearly; and in consequence, the triangle ABC exceeds the semicircle ADB by one thousandth part. It is upon such a singular coincidence as this that the following simple rule for approximate rectification of the circle with compass and ruler has been framed.

Rule. — “From an external point, at a distance equal to the radius from the given circle, describe a circle cutting the given circle at the two extremities of that diameter which is perpendicular to the line joining the point with the centre of the circle; and perpendicular to the same line draw two tangents to the two circles, the first touching the original circle at the point most remote from the external point, and the second touching the other circle, and cutting the original circle in two places. Then the two chords joining these two last points with the point of the given circle nearest to the external point, will represent the two quadrants that are there terminated; and if these chords be produced until they meet the tangent first

drawn, they will intercept upon it a portion bisected at the point of contact, and representing the two quadrants that are there terminated: and the triangle which this tangent forms with the two chords so produced will represent the area of the circle: and each of these rectilinear quantities will exceed the corresponding curvilinear quantity which it approximately represents by one thousandth part of the whole, nearly."

In proof of this construction, it will be seen that both DG and EK (Fig. 2) have by the construction the numerical values

Fig. 2.



$\sqrt{[2 \{ \sqrt{5} - 1 \}]}$, radius being unity, and are therefore equal to one another, whence the inference proceeds as before.

This construction closely rivals in descriptive geometry the well-known arithmetical approximation to the value of π that is found in the vulgar fraction $\frac{22}{7}$, this fraction exceeding the true value by only four, and the above geometrical result exceeding the same by only twelve parts in ten thousand. This simple process will, for example, yield for the quadrant of a circle ten inches in diameter, a result exceeding the truth by less than the hundredth part of an inch, which amply suffices in accuracy for all that such processes can have in view.

P. S. — It has been pointed out to me that the equation $\sec x = \cot x$, which the angle $\tan^{-1} \frac{1}{2} \pi$ so nearly solves, is equivalent to that proposition which Herodotus has recorded concerning the dimensions of the great pyramid of Gizeh, when he says that the area of the sloping side is equal to the square upon the perpendicular height. The semiangle contained at the vertex between two opposite sloping sides is in this case such an angle as x , and

accordingly the above construction may be considered as affording a ready means for reproducing upon paper those proportions which a recent writer* has affirmed that this monument was intended to perpetuate, the periphery of its base bearing to its height very nearly the proportion of the circumference of a circle to its radius. The triangle AKL in the above construction will accordingly represent the vertical section of the Great Pyramid parallel to its sides. — A. S. HERSCHEL, Collingwood, England, August 18, 1860.

Mathematical Monthly Notices.

General Problems from the Orthographic Projections of Descriptive Geometry; with their Applications to Oblique — including Isometrical — Projections, Graphical Constructions in Spherical Trigonometry, Topographical Projection ("One Plane Descriptive"), and Graphic Transformations. By S. EDWARD WARREN, C. E., Professor of Descriptive Geometry and Geometrical Drawing in the Rensselaer Polytechnic Institute, Troy, N. Y. New York: John Wiley, 56 Walker Street, 1860. 8vo. pp. xxxv, 412.

The aims of PROFESSOR WARREN in this work are thus stated in his Preface: —

"1st. To proceed upon the basis of a perfectly accurate and rational idea of what Descriptive Geometry is.

"2d. To make the course perfectly *determinate* and rational, even to the very details, in the plan of its composition and the order of its progression.

"3d. To embody in its methods of treatment, as well as in its preparatory directions to the student, safeguards against false modes of study and sham scholarship.

"4th. To make the course sufficiently full to meet the wants of advanced students, and of the members of the professions based on mathematics and the physico-mathematical sciences."

The plan of the course is founded on the division of Surfaces into Ruled Surfaces and Double-curved Surfaces, and the subdivision of Ruled Surfaces into Plane Surfaces, Single-curved Surfaces, and Warped Surfaces. Upon each of these four classes of surfaces four operations are distinguished, namely, Projections, Intersections, Tangencies, and Developments.

In carrying out this plan the *General Problems* are placed under the two heads of *Lower Descriptive Geometry* and *Higher Descriptive Geometry*. Under the first head are treated quite fully the various problems relating to the plane, straight line, and point; those relating to the cone and cylinder, including conic sections; those relating to the hyperboloid of one nappe, and the hyperbolic paraboloid; and lastly, those relating to the sphere, ellipsoid, elliptic paraboloid, and hyperboloid of two nappes. Under the second head we find the construction of various spirals, the cissoid, conchoid, involutes, cycloids, helix, and spherical epicycloid; also problems on the developable helicoid, warped oblique arch, oblique helicoid, right conoid, and annular torus.

* "The Great Pyramid, and why it was built," by JOHN TAYLOR.

Book II. gives the *Applications of Orthographic Projections*. These include Isometrical Projections, Conventional Oblique Projections, Spherical Trigonometry, Topographical Projections (a name which the author prefers to One-plane Descriptive Geometry), and Graphic Transformations, which include changing the planes of projection and rotating given magnitudes. A collection of Miscellaneous Problems, sixty-three in number, closes the work.

From the above it will be seen that PROFESSOR WARREN'S book is far more comprehensive than any work on Descriptive Geometry before offered to American, or, so far as we know, to English readers. As such it will doubtless be welcomed by the Professors in our Scientific Schools, and by many students to whom the extensive French works on this subject are inaccessible. But, besides its comprehensiveness, this book has many positive merits. The author appears to have studied his subject well, and to be master of its minutest details. He is without doubt familiar with the works of his predecessors; but the marks of independent thought are apparent on even a cursory examination. His experience as a teacher has enabled him to judge of the wants of students, and to make many useful suggestions in the shape of remarks. We find also several examples of extended discussions of problems, that show a praiseworthy love of thoroughness.

With this general recognition of the merits of the book before us, we pass to some more particular remarks upon a few points.

The true sphere of Descriptive Geometry is one of the topics discussed in the Preface, and the conclusion arrived at is, that Descriptive Geometry is "the *exact graphic art*, and not the science of geometry, and is distinctively, *graphically operative geometry*, or the *Geometry of Graphic Problems*." In this the author agrees with CHARLES (LIOUVILLE'S *Journal*, 1847, p. 33), who also denies to Descriptive Geometry the name of a science. But there are many able supporters of the opposite view, among whom may be noticed MONGE, CARNOT, PONCELET, and OLIVIER. M. OLIVIER discusses the subject quite fully in the preface to his *Additions au Cours de Géométrie Descriptive*, and becomes quite indignant against those who deny the claims of Descriptive Geometry to be called a science as well as an art.

PROFESSOR WARREN has also some remarks on the use of models and pictorial diagrams in teaching Descriptive Geometry, and condemns their general use, on the ground that without them certain powers of the mind — attention, abstraction, and especially conception — are better cultivated. This objection would be more forcible, if these powers could be cultivated in no other way, or if there were not many persons to whom Descriptive Geometry would be very useful, if it could be brought within their grasp without long and to them painful study. The truth is that no real simplification of any science is to be dreaded, even when we regard study as a discipline of the mind. When fields once inaccessible to ordinary minds are annexed to the common domain, there will always be higher fields opening up to exercise and strengthen the loftiest faculties.

OLIVIER, with an experience of more than twenty years in teaching Descriptive Geometry, recommends the use of models in the strongest terms. The costly models bearing his name are well known; but he has also contrived a simple apparatus which any one can have made for himself, and which we shall here describe for the benefit of those who have not access to this author's works. Take two shallow boxes similar to the tables of a backgammon board, and connected like them by hinges. Fill up these boxes with sheets of cork, and attach a brace to hold one table at right angles to the other, when necessary. These are the planes of projection. Provide four sets of rods from one eighth to one fifth of an inch square, and of various lengths from four inches to twelve, or sixteen, or twenty four inches, or even longer, according to the size of the planes. These rods are to be pointed with needles, that they may be readily fixed in any position in the cork. They are to be painted, one set red, one red and white in

bands, one black, and one black and white in bands. The red rods are used to construct in space any system referred in position to the two planes of projection; the red and white rods to project the points of this system on each of the planes of projection; the black and white to project these projections on the ground line; and the black to represent the projections of the red lines in space. When by means of these rods the operations necessary for the solution of a problem have been completed, the red and the red-and-white rods are removed, and the vertical plane revolved on its hinges to a horizontal position. The work is then done. By this apparatus also all ruled surfaces, such as cones, cylinders, hyperboloids of one nappe, hyperbolic paraboloids, conoids, &c., may be exhibited.

The classification of subjects adopted by PROFESSOR WARREN will arrest the attention of every one. Few, we think, will approve it. System is in itself a great good; but here the dividing, subdividing, and dividing again, are carried so far, that one almost doubts whether the confusion incident to so many heads is not greater than that of no classification whatever. A few distinct heads, such as those generally adopted by writers on this subject, are all that are necessary, at least in a text-book.

The style is also frequently open to objection. The greatest simplicity, directness, and precision should characterize such a work. All long and wordy sentences, all parade and circumlocution, all vagueness and ambiguity, are serious hindrances to the student. As an example of prolixity (one of the worst perhaps), we make an extract from the chapter on Graphic Transformations.

"426. Beneath all the formal problems of Descriptive Geometry, and giving character to their mode of solution, is a tacitly recognized problem, founded in the failure of human inventive skill to keep pace with human knowledge, as seen in the incapacity, hitherto apparent, to make *available*, that is, *cheap* and *permanently reliable*, instruments for the *easy* and *exact* graphical construction, by a continuous movement, of any lines but right lines and circles, — other lines being usually determined by points which are afterward joined by hand, or by the use of the 'irregular curve' (ruler).

"427. In continuation of this reasoning, it appears: 1st. That any point revolves about a fixed axis in a circular arc; 2d. In order to secure an adequate, that is, an immediately available, representation of such an arc, it must appear circular or rectilinear in projection; 3d. For this purpose, the plane of the arc must be parallel or perpendicular to the plane of projection; and 4th. This being the case, the axis of the arc will be respectively perpendicular or parallel to the same plane of projection."

We understand by this simply, that, since straight lines and circles are more easily drawn than other lines, it is best to choose an axis of rotation either parallel or perpendicular to the plane of projection; because then the path of a rotating point will be in projection a straight line, or an arc of a circle.

On page 5 we find: "Any magnitude, except a point, which is irreducible to anything more simple, may be regarded as the result of the motion of some simpler magnitude, according to a certain law." Here it is twice implied that a point is a magnitude; while on page 1 it is said that, "Magnitudes are definite portions of *space*, having one, two, or three dimensions." A point is therefore a definite *portion of space*, and has at least one dimension, — probably three, since its length, breadth, and thickness are very likely equal.

On page 104 we find: — "All sections of an oblique cone with a circular base which are parallel to the base, are circles, whose diameter depends on their distance from the vertex. All other sections, whose planes cut all the elements, are ellipses, save the sub-contrary circular section, *EF*, Pl. VIII., Fig. 53, whose plane makes an angle with an element, as *VB*, equal to the angle made by the parallel plane, *CD*, with the opposite element, *VA*."

Here are two omissions. It should have been stated, that the angles with VB and VA are both to be taken on the side towards the vertex. Moreover, there is no intimation that the figure does not represent *any* section of the cone through opposite elements; while it is essential that it should represent the particular section made by a plane drawn through the axis perpendicular to the plane of the base.

In drawing tangents to curves, the author appears to have had in view the method of ROBERVAL. This method is thus stated by its author (*Mém. Acad. des Sciences*, VI. p. 22): —

“*Axiom.* The direction of the motion of a point which describes a curve is a tangent of the curve in each position of this point.

“*General Rule.* By the specific properties of the curve (which are given) examine the different motions which the describing point has at the place where you wish to draw the tangent: compound these motions into one, and draw the line of direction of the compound motion; you will have the tangent of the curve.”

PROFESSOR WARREN says (p. 72): —

“The tangent to a curve is therefore the direction in which the generatrix, when at the point of tangency, would move were it to cease to generate a curve, and actually to move in the direction of its tendency for the instant.

“But the law of the generation of a curve is such that the tendency of the generatrix at any instant is the resultant of two component tendencies, while from any common school philosophy we know that such a resultant corresponds to the diagonal of a parallelogram, whose two adjacent sides express the relative values and directions of the component tendencies.”

Then follow applications to the conic sections. On pages 284, 385 (the subject being higher curves) we find the method restated thus: —

“The tangent line to the curve, at any point, is the resultant direction of the motion of the generatrix, due to the two component motions of that generatrix.

“The usual case is that neither of these motions is uniform. In this case, the tangent at any point is the resultant of the two tendencies for that instant only. Hence to construct the tangent at any point, draw two straight lines through that point in the directions in which the generatrix tends to move, at that point, and lay off on these lines distances whose ratio shall equal the *ratio* of the *rates* at which the generatrix tends to move at the instant in the given directions. The diagonal of the parallelogram, formed on these distances as two of its sides, will be the required tangent.

“To furnish such momentary ratios as are necessary to the above solutions is one object of the higher calculus.”

Now the principle of ROBERVAL's method is acknowledged to be correct; but in applying it there are undoubtedly difficulties in determining what the components are. ROBERVAL himself, MONTUCLA, and even MONGE, in one case, are said by DUHAMEL (*Mém. Sav. Étrang.*, V. 1838, p. 257) to have confounded the component velocities of the generatrix with the projections of the total velocity upon the directions of the components. He shows also, that this confusion will produce no actual error, when the velocities are equal, or when the directions of the motions are at right angles to each other. But when a curve is given by an equation between the distances of its points to two or more foci, these conditions do not in general exist, and the error becomes manifest. To illustrate, suppose we had the equation $ar + br_1 = a$ constant, r and r_1 being the distances from any point of the curve to two fixed points, and a and b any multipliers, and let ϕ and ϕ_1 be the angles made by the tangent to the curve at any point with r and r_1 . Then it is shown by DUHAMEL and by SERRET (*Méthodes en Géométrie*, p. 53, *et seq.*) that

$$\frac{dr}{dr_1} = -\frac{b}{a} = \frac{\cos s}{\cos s_1};$$

and, consequently, as will be readily seen by drawing a figure, that if we lay off on r and r_1 any distances having the ratio $dr:dr_1$, and erect perpendiculars at the extremities of these distances, the intersection of the perpendiculars will give the direction of the required tangent. It is obvious that the direction will not be that of the diagonal of a parallelogram constructed on the distances laid off as above, unless either $dr = \pm dr_1$, or the angle between r and r_1 is a right angle.

When, therefore, this method is used, the mode of finding the ratio of the components should be distinctly laid down. We have seen that it is possible to fall into error, even when the Differential Calculus is relied on to furnish the ratios. At any rate, if we have to rely on the Calculus for this essential step, the method ceases to be an independent one, and we had better adopt at once the recognized methods of drawing tangents derived from Analytical Geometry and the Calculus.

Extended as these remarks have become, we have left unnoticed many points that deserve attention. Enough, however, has been said to call attention to the principal features of this work. While we have not hesitated to point out what we consider faults, we trust that we properly appreciate the labors of the author in giving to the English student so extended, and, in many respects, so useful a work.

The Lady's and Gentleman's Diary, or Poetical and Mathematical Almanack for the Year 1861. London: Price one shilling and six pence.

This is the 158th annual number of this interesting and important publication. It is impossible to estimate the influence this unpretending serial has exerted upon the progress of the mathematics, and especially upon the development of mathematical talent in England. Besides 1988 original problems with their solutions, embracing all departments of the science, the mathematical papers thus far published would make several most valuable volumes. In our next we shall give the problems proposed for the next year, in hopes that many of our younger readers will feel disposed to send solutions of them "to the Editor of the Lady's and Gentleman's Diary, Stationer's Hall, London," post-paid, before May 1st, 1861.

Elements of the Differential and Integral Calculus. By WILLIAM SMYTH, A. M., Professor of Mathematics in Bowdoin College. Second Edition. Portland: Published by Sanborn and Carter. 1859.

The luminous statement, given in the Preface, of the spirit and plan of the work, at once arrests the attention; and an examination shows that the author is not only master of his subject, but clearly comprehends the manner of its presentation for the purpose of instruction. We have no hesitation in saying that this is one of the very best text-books upon the subject we have ever seen. The infinitesimal method of LEIBNITZ is adopted as its basis, which the author considers less logical than NEWTON's Method of Limits or LAGRANGE's Method of Derived Functions, but better adapted to the purposes of instruction, both on account of its greater simplicity, and the facility with which it is applied to the solution of the ordinary problems demanding the calculus, and especially for all purposes of investigation. Differentials are regarded simply as auxiliary quantities, and those who are familiar with the views of CARNOT (to whom our author acknowledges his indebtedness), as exhibited in his incomparable little work, entitled, *Reflexions sur la Méaphysique du Calcul Infinitésimal*, will at once understand the spirit of the work before us. The arrangement is excellent, as the following brief synopsis will show. I. First Principles. II. Derivation of the auxiliary quantities, or differentials. III. Application to problems in which the differentials can be elim-

inated by the ordinary processes of Algebra. IV. Integral Calculus, in which the auxiliaries, or differentials, are eliminated by reversing the process by which they were obtained. V. Application to problems of quadrature, cubature, rectification; surfaces of solids. VI. Successive differentials, or new auxiliaries. VII. Application of calculus to development of functions. VIII. Maxima and minima. IX. Theory of curves. X. Transcendental functions, their differentiation and integration. XI. Transcendental curves. XII. Processes of integration. XIII. Cycloid. XIV. Quadratures and cubatures continued. XV. Application to Mechanics. XVI. Equilibrium, Centre of Gravity, Hydrodynamics. XVII. Method of Variations. XVIII. Application of Calculus to Astronomy. XIX. Limits, Derived Functions. XX. Miscellaneous examples.

The clearness and simplicity with which the principles are developed, the interesting historical references, and the variety of the problems, all combine to make the work before us an excellent text-book, and as such we commend it to teachers and students, with the assurance that they will not be disappointed.

The methods of NEWTON and LAGRANGE are very properly given at the end of the work, because the student is then prepared to compare them with the method of LEIBNITZ without confusion.

Editorial Items.

We are happy to lay before our readers the following report; and especially so, because all the Essays submitted are found of "sufficient merit to be worthy of publication." We shall print the Essays in the order named as rapidly as possible, and all in this volume of the Monthly.

REPORT OF THE COMMITTEE ON PRIZE ESSAYS.

The Committee on Prize Essays regret that their Report has been so long delayed. The protracted absence of one of their number and the engagements of the rest prevented an earlier decision. They have received four Essays: On the Conformation of the Earth; On Central Forces; On Projections; and On Spherical Conics. All of these they consider of "sufficient merit to be worthy of publication;" although, as the productions of students, they are not to be too severely criticised. The Committee award the following prizes: To the first Essay, in the order named, forty dollars; to the second, thirty dollars; to the third, twenty dollars; and to the fourth, ten dollars.

WILLIAM FERREL,
J. B. HENCK,
CHAUNCEY WRIGHT.

BOOKS RECEIVED. — General Problems from the Orthographic Projections of Descriptive Geometry, &c., by PROF. S. EDWARD WARREN. (See Notice.) For sale by Brown and Taggard, Boston. — Annual Report of the Board of Regents of the Smithsonian Institution, showing the operations, expenditures, and condition of the Institution for the year 1859. — On the Equation of Differences for an Equation of any Order, and in particular for the Equations of the Orders Two, Three, Four, and Five, by ARTHUR CAYLEY, F. R. S.; also a paper on Recent Terminology in the Mathematics. — Progressive Intellectual, Practical, and Higher Arithmetics; New Elementary and University Algebras; New Plane and Solid Geometry, with Plane and Spherical Trigonometry; Surveying and Navigation, by H. N. ROBINSON, LL. D., published by Ivison, Phinney, & Co., 48 and 50 Walker Street, New York.










Fig. 9.




Fig. 10.




Fig. 11.




Fig. 13.




Fig. 12.




Fig. 14.




Fig. 15.




Fig. 16.




Fig. 17.




Fig. 18.




Fig. 19



Fig. 20




Fig. 21




Fig. 22




Fig. 24





Fig. 23






Fig. 26.




Fig. 25.




Fig. 27.




Fig. 28.






Fig. 29.





Worcester's Quarto Dictionary *The Standard*

VERY SIGNIFICANT FACTS.

The following recommendations are from some of the most distinguished American and English scholars. They are but a few from many which have been received, testifying to the superiority of Worcester's Quarto Dictionary. These testimonials are of the highest value, for they have all been given during the present year, and after an examination of this work and of that which is endeavoring to hold the position of a rival. The scholars of America and of England, with scarcely an exception, have decided in favor of Worcester. Not a single scholar, equal in authority to any one mentioned below, can be cited as giving, after a comparison of the two works, the preference to Webster's Dictionary. We give the testimony:—

From C. C. FELTON, LL.D., President of Harvard College.

Aware of the labor and care which had been devoted to this (the department of scientific terms) as well as to other parts of the work, I felt assured that Worcester's Quarto Dictionary would more nearly meet the public wants than any other hitherto published.

My expectations have been more than fulfilled. I find it not only rich beyond example in its vocabulary, but carefully elaborate in all the details, and thoroughly trustworthy as a guide to the most correct and elegant usage of the language.

From the REV. JOSEPH BOSWORTH, D.D., F.R.S., Professor of Anglo-Saxon, Oxford, England.

It is the most complete and practical, the very best, as well as the cheapest English Dictionary that I know.

From GEORGE P. MARSH, LL.D.

The work of Dr. Worcester is unquestionably much superior to any other general dictionary of the language in every one of these particulars (orthography, pronunciation, definition, fullness of vocabulary, and precision and distinctness of definition).

From REV. W. WHEWELL, D.D., Master of Trinity College, England.

I have repeatedly consulted the Dictionary since it has been in my possession, and have seen reason to think it more complete and exact than its predecessors.

From CHARLES RICHARDSON, LL.D., the oldest living English Lexicographer, England.

I sincerely hope you may enjoy from your brethren, both in America and England, that tribute of honor to which you have earned so undoubted a title.

From D. R. GOODWIN, D.D., President of Trinity College, Hartford.

It was but a short time since that I was led to commend another dictionary as, on the whole, and with some exceptions, the best and most complete thing of the kind within my knowledge. The commendation was honestly given at the time; but now it must be withdrawn in favor of yours. I consider your dictionary, in orthography, pronunciation, and definitions, as superior to any of its predecessors.

From REV. W. B. SPRAGUE, D.D., of Albany, N. Y.

My opinion of Worcester's Quarto Dictionary, after having given it as extended an examination as my circumstances would admit, is, that there is no other dictionary in the language that compares with it for completeness, accuracy, comprehensiveness, and precision, and perhaps I ought to add, that I have arrived at this conclusion rather contrary to a preconceived opinion.

From REV. HENRY A. BOARDMAN, D.D., of Philadelphia.

I particularly like it (the Dictionary). 1. Because of its very comprehensive character: 2. Because it adheres to the settled orthography of our noble language,—discarding those innovations which, however countenanced by certain publishing-houses, have never to any extent been accepted by the scholars of our country.

From LOUIS AGASSIZ, LL.D.

It is of great importance, when the nomenclature of science is gradually creeping into common use, that an English lexicon should embrace as much of it as is consistent with the language we speak. I am truly surprised and highly delighted to find you have succeeded far beyond my expectations in making the proper selection, and combining with it a remarkable degree of accuracy. More could hardly be given except in a scientific cyclopædia.

The following lines are quoted from Harper's Magazine for September. They serve to show very truthfully the comparative value of recent and old commendations:—

"INJUSTICE.—Our attention has been called to a species of injustice of which publishers are sometimes guilty, in publishing commendations of school-books, without giving the dates when they were written. Especially does this merit reproof when these commendations are old, and when it is known that the writers have subsequently commended other and later publications in the same department. It will readily be seen that this is frequently not only an act of injustice to teachers who have had the courtesy to commend a book, but that it is also a fraud upon the public."

SWAN, BREWER, & TILESTON,

131 Washington Street, Boston.

No. 117 Washington Street, Boston,
October, 1860.

CROSBY, NICHOLS, LEE & CO.

HAVE JUST PUBLISHED

THE WORKS OF CHARLES LAMB. With a fine Portrait. In Four Volumes. 12mo. Price, \$5.00.

This new edition of Lamb is reprinted from the latest London edition, edited by Talfourd, contains his Life, Letters, and Final Memorials, his Essays of Elia, Rosamund Gray, with other Essays and Poems, being the best and most complete edition issued.

It is elegantly printed on paper slightly tinted, and is in every respect an edition worthy of the author.

"This is the most convenient and elegant edition of Lamb's Works which has been published, either in England or in this country, and it leaves nothing to be desired in respect to typographical beauty or any other essential of a good library edition."—*North American Review*.

The Three Great Histories,—Hume, Gibbon, and Macaulay.

Elegant Library Editions, beautifully printed on tinted paper, convenient in size, and withal the best and neatest yet published.

THE HISTORY OF ENGLAND, FROM THE INVASION OF JULIUS CÆSAR TO THE ABDICATION OF JAMES THE SECOND, 1688. By DAVID HUME, Esq. A new Edition; with the Author's last Corrections and Improvements. To which is prefixed a short Account of his Life, written by himself. In 6 volumes. 12mo. With a Portrait. Price, \$7.50.

SIR JAMES MACKINTOSH says of Hume: "His greatest work, and that which claims most attention, was his 'History of England,' which, notwithstanding great defects, will probably be at last placed at the head of historical compositions. No other narrative seems to unite, in the same degree, the two qualities of being instructive and affecting. No historian approached him in the union of the talent of painting pathetic scenes with that of exhibiting comprehensive views of human affairs. . . . His narrative is flowing and various; in common events, short and clear; in great actions, rapid and animated; in affecting incidents, circumstantial and picturesque. His general observations seem always to be required by his subject; the most profound ideas are clothed in a transparent simplicity."

THE HISTORY OF THE DECLINE AND FALL OF THE ROMAN EMPIRE. By EDWARD GIBBON, Esq. With Notes by the Rev. H. H. MILMAN. A new edition. To which is added a complete Index of the whole Work. In 6 volumes. 12mo. With a Portrait. Price, \$7.50.

Of the various editions of GIBBON this of MILMAN is uniformly acknowledged to have no rival.

PROF. SMYTH, in his *Lectures on Modern History*, says: "But the whole is notwithstanding, such an assemblage of merits, so various, so interesting, and so rare, that the History of the Decline and Fall must always be considered as one of the most extraordinary monuments that have appeared of the literary powers of a single mind, and its fame can perish only with the civilization of the world."

THE HISTORY OF ENGLAND FROM THE ACCESSION OF JAMES II. By THOMAS BABINGTON MACAULAY. In 4 volumes. 12mo. Price, \$5.00.

"It is as fluent and as much colored as Livy; as close and coherent as Thucydides; with far more real condensation and a larger thoughtfulness than either; and quite free from the affected laconisms and sarcasms and epigrams of Tacitus. I do not know that I ever read anything so good as the first forty pages; so clear, comprehensive, and concise, so pregnant with deep thought, so suggestive of great views, and grand and memorable distinctions."—LORD JEFFREY.

CURIOSITIES OF LITERATURE. By ISAAC DISRAELI. With a View of the Life and Writings of the Author, by his Son. In 4 volumes. 12mo. Price, \$5.00.

"It embraces a more copious selection of noteworthy facts in regard to books and authors than can be found in almost any other compilation of the kind, while it doubtless owes much of its popularity with all classes of readers to the agreeable form in which these facts are presented."—*North American Review*, April, 1860.

THE NORTH AMERICAN REVIEW. No. CLXXXIX.—For October, 1860.


CONTENTS.—Article I. Homer and his Heroines. II. Climatology. III. Life and Labors of Thomas Prince. IV. Edmund Waller. V. Lord Shaftesbury. VI. Second Volume of Palfrey's "History of New England." VII. Quarantine and Hygiene. VIII. Rush's Occasional Productions. IX. The English Language in America. X. The Origin of Species. XI. An "Inglorious Milton." XII. Critical Notices. XIII. New Publications.

This Review has for nearly fifty years stood at the head of the *litera* our country, and has numbered among its contributors most of our literary men. The writings of Webster, Everett, Bancroft, and were first made known through this work; and its pages are now articles from our most prominent writers.

The NORTH AMERICAN REVIEW is published quarterly, in numbers of three hundred pages each, at five dollars a year.

CHILSON'S PATENT CONE FURNACE.

PATENTED IN
AMERICA, ENGLAND, AND FRANCE.



The subscriber begs to call the attention of scientific gentlemen and others to his discovery of a principle for burning fuel, by which all the combustible properties of the smoke and gases eliminated from the burning fuel are thoroughly consumed and their heat retained and made available to the fullest extent, so that there shall be no waste whatever. This important object is accomplished by the use of a large cluster of cones or tapering radiators placed over a broad, shallow, flaring fire-pot, and surrounded by a radiating and exhaustive annular chamber, as represented in the cut. The cones being large at their base and small

at the tops, causes the smoke and gases to be suspended directly over and near the fire, where they are thoroughly consumed, and their heat forced to impinge against and up the cones, thereby creating a powerful radiation.

This plan of furnace secures a fresh, healthful heat, free from that scorching, dry, disagreeable, red-hot iron heat and coal-gas so common to hot-air furnaces generally. Persons interested in these matters will here find an invention worthy of their attention, which saves from one half to two thirds of the fuel usually consumed.

COOKING RANGES.

CHILSON'S PATENT DOUBLE OVEN RANGE.

The most important Improvements yet made in Ranges.

THE highly satisfactory testimony received from the large number of these Ranges in use in the past three years should certainly induce those in want to examine this Range, which is *entirely new in principle*, and different from any other before known. It has been got up very perfect and substantial so that it commends itself at once to the purchaser, in preference to anything else of the Range kind in this country.

Its great economy in the use of fuel, its quick and even baking ovens, its nice arrangement of flues, — without the usual vexatious complication of dampers, — its cast-iron flue plates, preventing any loss of heat in the brick-work, its nice arrangement for roasting meats in the ventilated ovens, &c., will be appreciated by every housekeeper. Sizes in variety for dwellings, hotels, &c., with or without Water Backs for heating water for bathing, &c., Hot Closets, and Hot Air Fixtures for heating additional rooms.

Also, a full assortment of

Chilson's Patent Trio Portable Furnaces, Trio Stoves, Parlor Stoves, Registers, Ventilators, &c., &c.

WAREROOMS 99 & 101 BLACKSTONE ST., BOSTON.
FOUNDRY AT MANSFIELD, MASS.

GARDNER CHILSON.



Philosophical Instruments

AND

SCHOOL APPARATUS,

MANUFACTURED BY

E. S. RITCHIE, BOSTON, MASS.

Few branches of study in our schools are of more practical importance than Natural Philosophy, and it is only by the use of proper apparatus that its principles can be adequately elucidated.

Until a comparatively recent period, the attention bestowed upon this subject has been confined to a few of the elementary principles; of late years, however, the growing interest in physical science, as a branch of common education, has created a desire for more ample and varied means of experimental demonstration.

It has been my earnest desire and effort to keep pace with the progress of natural science, and, as new principles have been discovered or developed, to devise and construct new instruments, to improve the old, and to add to my list many which heretofore could only be procured from Europe; and also that, in superior quality and finish, as well as in graceful form and proportions, whatever goes from my manufactory shall take rank with the best products of European make.

So numerous and important are the changes that have consequently been made, that I have published a new and enlarged Catalogue of 100 pages octavo, containing descriptions of over

Twelve Hundred Pieces of Apparatus,

and illustrated by over **Three Hundred** finely executed engravings, and also many commendatory letters from the most distinguished Physicists in the country.

E. S. RITCHIE.

Philosophical Rooms, No. 313 Washington Street, Boston.

☐ Catalogues will be sent, prepaid, by mail to order, — please enclose 25 cents in stamps.

PHOTOGRAPHING ON WOOD.

BRICHER AND RUSSELL,

Photographers, Designers, and Engravers
ON WOOD.

129 WASHINGTON STREET, BOSTON.

BRICHER AND RUSSELL, by their new process of Photographing on Wood, are now enabled to produce much more accurate representations of objects than by the old and slow process of hand drawing. They will furnish Portraits, Views of Colleges, Seminaries, Academies, and Public Buildings generally, College and Society Seals, Philosophical Apparatus, Mathematical Instruments, and Diagrams (see specimens in Mathematical Monthly), Illustrations for School Books of all kinds, &c., &c.

Directions.

Get a Photographer to take a *negative* of the object you wish to have engraved, and of the required size, — that is, send us a photograph of the required size, and the *reverse* of what is wanted when printed.

If a negative photograph cannot be obtained, get as sharp and clear an ambrotype as possible.

ROBINSON'S COURSE OF MATHEMATICS.

BY HORATIO N. ROBINSON, LL.D.,

Late Professor of Mathematics in the United States Navy.

The most Complete, most Practical, and most Scientific Series of Mathematical Text-Books for Common Schools, Academies, and Colleges ever issued in this country.

NUMBERING EIGHTEEN VOLUMES

Of well-graded and progressive text-books, carefully and thoroughly prepared, in which the Author has been assisted by some of the best practical Mathematical talent in the country.

The present edition of the *Analytical Geometry and Calculus*, in one volume, while it contains very much that is valuable and is much liked as a text-book, yet doubtless it is liable to criticisms and may be improved. For this reason the work is being *re-written and thoroughly revised*, in which is engaged some of the best mathematical talent in the country, and will appear early in the Spring of 1861 in two handsome octavo volumes.

Robinson's New Geometry and Trigonometry, recently published, is a beautiful volume, and worthy the attention of the thorough and practical teacher, who desires a first-class text-book on those subjects.

The publishers have received an immense number of commendations for the books of this Series, of which the following are selections:—

Professor W. H. SEAVEY, A. M., Principal of Girls' High and Normal School, Boston:—"I regard Robinson's Universal Astronomy as the best work on the subject, within the range it professes for itself, that I have ever seen. It is not a compilation. If everything in it is not *n-w*,—of course it cannot be in such a work,—everything in it is presented with the same living freshness and spirit that constitute a peculiar excellence in Professor Robinson's other educational works, particularly his Algebra and Geometry. To the printed page he has imparted the power of awakening an interest and zeal scarcely inferior to that aroused by the living voice of the enthusiastic teacher. Professor Robinson's books are not labor-saving machines: they invite to labor. It is no part of their object to keep difficulties out of sight, but to present them in such a manner as to inspire the student with confidence in his own ability to cope with them, and an ardent desire to encounter them."

Prof. J. W. PATTERSON, A. M., Professor of Mathematics, Dartmouth College:—"Robinson's Works possess superior merits."

Prof. S. S. GREENE, A. M., Brown University: "Robinson's Mathematics are an excellent series."

Prof. H. POMEROY, C. E., Lawrence University, Wis.:—"Robinson's Mathematical Operations," in connection with the sections upon logarithmic calculations in his 'Navigation,' are far finer than anything else of the sort I know of in our language. Even in the elaborate work of Babinet and Housel, just published in Paris (*Calculs Pratiques*), there is nothing equal to the above named sections on the niceties of logarithmic computation. They contain a mass of matter useful in the office of the Engineer, in the camp of the Scientific Explorer, and in the fixed observatory of the Astronomer."

F. G. CAREY, late President of Farmers' College, Ohio:—"You ask my opinion in relation to Prof. Robinson's Mathematical Works. I reply, that the best evidence of our appreciation of them is, that we have adopted them throughout as text-books in this College, and have found by experience that they contain just what such text-books should contain, and show their author to be not only a book-maker but a practical teacher."

Prof. J. O. HUDNUT, C. E., of Chicago:—"In a long career of travel which ten years' experience as an engineer has given me. I have never met Professor Robinson's equal as a practical mathematician. His books are unrivalled for perspicuity and clearness in our language. His University Algebra and Astronomy are both jewels in the mines of science. Take the course together, I believe it to be the best in the English language. I have the opinion of several practical engineers upon his Trigonometry, and they agree

with me in thinking it the most excellent treatise upon that subject yet published."

HON. JONATHAN TENNEY, Secretary Board of Education in New Hampshire, writes:—"I believe your Algebra to be the best yet published for the use of the students in the High Schools and Colleges of our country. The Geometry I shall introduce when we have a class, which will probably be next term."


Prof. W. D. WILSON, Hobart Free College, Geneva, N. Y., author of Treatise on Logic:—"I have obtained a copy of your Analytical Geometry and Calculus, and am exceedingly pleased with it. I regard it as incomparably more easy and intelligible for the beginner than any other work I have ever seen."

Prof. DASCOT GREENE, A. M., Professor of Mathematics, Polytechnic Institute, Troy, N. Y.:—"Robinson's University Algebra is a very full and, at the same time, concise work. It is an *original* work which is more than can be said of most text-books on that science. The Geometry is, in many respects, an improvement on most works of its class. The Analytical Geometry and Calculus abound with that variety of illustration, and fulness of practical application so characteristic of the author. The treatise on Surveying and Navigation is an excellent practical work. The Mathematical Operations is not only valuable as a key to the former volumes of the series, but also as containing additional problems, involving many interesting processes. The Astronomy is a very excellent work, and contains more of physical astronomy than is often found in an elementary treatise. It is remarkable what an amount of astronomical science, descriptive, physical, and practical, is here compressed."

Prof. D. WOOD, A. M., Professor of Civil Engineering and Physics in the University of Michigan:—"For conciseness, accuracy, and adaptation to classes I think Robinson's Mathematical Series unequalled by any in this country."

Prof. JAMES C. WATSON, A. M., Director of Observatory, University of Michigan:—"I consider Robinson's Mathematical text-books the very best yet published in this country. Their introduction ought to become general."

Prof. T. McCauley Ballentine, A. M., Professor of Mathematics, Cumberland College, Ky.:—"Robinson's Analytical Geometry and Calculus will be found very valuable books. The development of the various subjects treated is so clear and natural, that the student who has had sufficient preparation can thoroughly understand them."

 A NEW DESCRIPTIVE CATALOGUE (of 160 pages), and the "EDUCATIONAL CIRCULAR," with matter interesting to Teachers, descriptions, notices, testimonials, prices, etc., sent free, on request.

 Liberal Terms for Specimen Copies and first supplies for Schools.

IVISON, PHINNEY, & CO.,

Publishers, 48 and 50 Walker Street, New York.

SOLD BY BOOKSELLERS GENERALLY EVERYWHERE.

ROSS' AMERICAN SCHOOL FURNITURE WORKS,

Established, 1838.

The Pioneer and most Extensive Establishment in the United States.



The above Engraving represents Ross' Improved High School, Normal School, or Academy Double-Column Desk and Chairs; Falls to lift; plain top.



The above Engraving represents Ross' Improved High School, Normal School, or Academy Single-Column Desk and Chair; Fall to lift; plain top.



The above Engraving represents Ross' Improved High School, Normal School, or Academy Single Desk and Chair; Fall to lift; plain top.



The above Engraving represents Ross' Improved High School, Normal School, or Academy Double Desk and Chairs; Falls to lift; plain top.

An Illustrated Catalogue and Information forwarded, on Application, by Mail or Otherwise.

N. B. *Every Article of School Furniture from this Establishment WILL BE WARRANTED. All Communications may be addressed to*

JOSEPH L. ROSS,
Office--CHARDON, opp. HAWKINS STREET,
(Near the Revere House,) BOSTON, Mass.

Entered according to Act of Congress, in the Year 1860, by JOSEPH L. ROSS, in the Clerk's Office of the District Court of Massachusetts.

MATHEMATICAL BOOKS,

IMPORTED AND FOR SALE BY

SEVER AND FRANCIS, CAMBRIDGE,

SUCCESSORS TO JOHN BARTLETT.

ENGLISH.

American Ephemeris and Nautical Almanac. For the Years 1855-61. Each	\$ 1.50
BAKER. Treatise on Statics and Dynamics	0.30
BLAND. Algebraical Problems. 8vo. Second-hand	1.50
Key to do. 8vo. do.	1.75
Philosophical Problems. 8vo. do.	1.50
Geometrical Problems. 8vo. do.	1.50
BOOLE. Treatise on Differential Equations	4.20
BOOTH. Theory of Elliptic Integrals. 8vo.	2.25
BORDEN. System of Railway Formulas. 8vo.	2.25
BRITISH NAUTICAL ALMANAC. 1861 and 1862, each	1.00
CARMICHAEL. Treatise on the Calculus of Operations. 8vo.	2.75
CHAUVENET. Trigonometry. 8vo.	1.50
COURTENAY. Calculus. 8vo.	2.50
COOMBE. Solutions of the Cambridge Problems, 1840, 1841. 8vo.	2.25
DEMORGAN. Differential and Integral Calculus	2.00
EARNSHAW. Dynamics. 8vo.	4.25
FISCHER. Logarithmic Tables of Seven Places. Translated from Bremiker's Vega. 8vo. Hf. mor.	2.50
FROST. Mathematical Questions of the Senate-House Examination Papers. 1838-1849. 8vo.	3.00
GAUSS. Theoria Motus. Translated by C. H. Davis. 4to.	5.00
GREGORY. Differential and Integral Calculus by Walton. 8vo.	5.30
HADDON. Differential Calculus	5.30
HAMILTON. Lectures on Quaternions. 8vo.	6.30
HANN. Integral Calculus	.30
HEATHER. Descriptive Geometry	.30
HEMMING. Differential and Integral Calculus. 8vo.	2.75
HUTTON. Mathematics	4.00
Recreations. 1 vol. 8vo.	4.00
Mathematical Tables. 8vo.	3.75
HYMER. Plane and Spherical Trigonometry	2.50
Algebraic Equations	3.00
Treatise on Conic Sections. 8vo.	2.75
Differential Equations and Calculus of Finite Differences. 8vo.	3.00
Integral Calculus	3.00
Geometry of Three Dimensions. 8vo.	3.00
JAMIESON. Solutions of the Senate-House Rider. 8vo.	2.25
LARDNER. System of Algebraic Geometry. 8vo. Second-hand	1.75
MAIN and BROWN. Indicator and Dynamometer	1.40
Mathematical Monthly. Edited by J. D. Runkle. Vol. I. In Nos. Do. Hf. mor.	3.00
Mathematical Problems and Examples of the Senate-House Examination Papers. 1821-1853. With an Appendix containing the Senate-House Questions for 1857. 8vo.	3.00
PARKINSON. Elementary Mechanics. Post 8vo.	2.87
PEIRCE. Analytic Mechanics. 4to.	7.50
PEIRCE, J. M. Analytic Geometry. 8vo.	1.50
PHEAR. Elementary Hydrostatics. Post 8vo.	1.02
PRICE. Differential and Integral Calculus. 3 vols. 8vo.	14.50
Quarterly Journal of Pure and Applied Mathematics. Sylvester, Ferris, & Co. \$6.00 per annum, post-paid.	
RANKINE. Manual of Applied Mechanics	2.00
SALMON. Conic Sections	3.50
Higher Plane Curves	3.50
Higher Algebra	1.80
Solutions of the Cambridge Problems. 1843-1851. Ferris and Jackson. 8vo.	4.50
1854. Walton and McKenzie. 8vo.	2.50
1857. Campion and Walton. 8vo.	2.50
Senate-House Riders. Jamieson. 8vo.	3.25
SPOTTISWOODE. Elementary Theorems relating to Determinants. 4to.	1.50
TATE. Exercises on Mechanics, with Key. 2 vols.	1.72
Mechanical Philosophy. 8vo.	3.75
Materials. 8vo.	1.75
TATE and STEEL. Dynamics. 8vo.	3.00
TODHUNTER. Analytical Statics. Post 8vo.	3.00
Differential and Integral Calculus. 2 vols. Post 8vo.	6.00
WALTON. Problems illustrative of Plane Coördinate Geometry. 8vo.	4.81
Mechanical Problems. 8vo.	5.50
Problems in Hydrostatics and Hydrodynamics. 8vo.	3.00
On the Differential Calculus. 8vo.	3.25
WALTON and MCKENZIE. Solutions of the Cambridge Problems. 1854. 8vo. \$3.25. Do. 1857.	2.50
Problems in Elementary Mechanics. 8vo.	3.00
WAUD. Algebraic Geometry. 8vo. Hf. cf. (Second-hand.)	1.50
WHEWELL. Analytic Statics. 8vo.	2.25
WILSON. Dynamics. 8vo.	2.87
WOOLHOUSE. Differential Calculus	.30

FRENCH, &c.

AUDÉ. Pousseée des Terres. 8vo. Hf. cf.	\$ 1.87
BELANGER. Résumé de Leçons de Géométrie Analytique et de Calcul Infinitésimal. 8vo.	2.75
BIOT. Traité des Equations différentielles. Hf. mor.	2.00
BOURCHARLAT. Eléments de Calcul Differential et du Calcul Integral. 8vo.	3.00
BONNET. Leçons de Mécanique Élémentaire à l'Usage des Candidats. L'Ecole Polytechnique. 8vo.	2.00
BRIOSCHIE. Théorie des Determinants. 8vo.	2.00
BOURDON. Application de l'Algèbre à la Géométrie. 8vo. Hf. cf.	3.00
CALLET. Tables de Logarithmes. 8vo. Hf. cf.	5.00
CHARLES. Traité de Géométrie Supérieure. 8vo. Hf. cf.	6.00
CHOQUET. Traité élémentaire d'Algèbre. 8vo. Hf. cf.	2.00
COURNOT. Traité élémentaire de la Théorie des Fonctions, et du Calcul Infinitésimal. 2 vols. 8vo.	4.00
CRELLE. Journal für die reine und angewandte Mathematik. Per annum	4.00
DELIBLE. Géométrie Analytique. 8vo. Hf. cf.	3.00
DUMAMEL. Cours d'Analyse Calcul Infinitésimal. 2 vols. 8vo. Hf. mor.	5.00
Cours de Mécanique. 3 vols. 8vo. Hf. mor.	4.50
DUPIN. Application de la Géométrie et de Mécanique à la Marine aux Ports et Chaussées. 4to. Hf. mor.	4.50
Developpements de Géométrie. 4to. Hf. mor.	4.50
Géométrie et Mécanique, appliquée aux Arts. 8vo. Hf. mor.	2.50
FRANCEUR. Cours complet de Mathématiques pures. 2 vols. 8vo.	5.00
Eléments de Statique. 8vo. Hf. cf.	2.00
GAUSS. Méthode des Moindres Carrés. 8vo. Hf. cf.	1.02
KÖHLER. Logarithmisch Trigonometrisches Handbuch. 8vo.	2.12
LACROIX. Traité du Calcul Differential et du Calcul Integral. 3 vols. in 4to, avec 18 pl. Mor.	35.00
LAGRANGE. Mécanique Analytique. 2 vols. 4to.	13.00
Théorie des Fonctions Analytiques. 4to. Hf. mor.	6.00
Traité des Equations Numériques de tous les Degrés. 4to. Hf. mor.	5.00
LALANDE. Tables de Logarithmes. 18mo.	1.25
LAPLACE. Système du Monde. 4to. Hf. mor.	5.50
LEGENDE. Traité des Fonctions Elliptiques. 3 vols. 4to. Hf. mor.	20.00
LEROY. Traité de Stéréotomie comprenant les Applications de la Géométrie Descriptive à la Théorie des Ombres, la Perspective Lineaire, la Gnomonique, la Coupe des Pierres, et la Charpente. In 4to, avec Atlas de 74 pl. in fol. Paper, \$9.00; Hf. mor.	12.00
Traité de Géométrie Descriptive. In 4to, avec Atlas de 71 pl. Paper, \$3.00; Hf. mor. \$5.00 and	6.00
Analyse Appliquée à la Géométrie des trois Dimensions. 8vo. Hf. cf.	2.25
LILOVILLE. Journal de Mathématiques Pures et Appliquées. 1836-1857. 22 vols.	
MABISTRE. Cours de Mécanique Appliquée. 8vo.	3.00
MONGE. Application de l'Analyse à la Géométrie. 4to. Hf. mor.	10.50
Traité élémentaire de Statique. 8vo. Hf. mor.	1.75
MORIN. Aide-mémoire de Mécanique Pratique. 8vo.	2.00
MONTFERRIER. Encyclopédie Mathématique ou Exposition complète de Toutes les Branches des Mathématiques d'après les Principes de la Philosophie des Mathématiques de Hoëné Wronski. 3 vols. 8vo. To be continued in monthly Livraisons.	6.00
NAVIER. (Ketten) Hangbrücken. 4to.	5.00
OLIVIER. Théorie Géométrique des Engrenages. 4to. Hf. mor.	3.75
Developpements de Géométrie Descriptive. 2 vols. 4to. Hf. mor.	9.00
Complément de Géométrie Descriptive. 2 vols. 4to.	9.00
Mémoires de Géométrie Descriptive, Théorique, et Appliquée. 2 vols. 4to. Hf. mor.	9.00
Application de la Géométrie Descriptive. 2 vols. 4to.	10.00
Cours de Géométrie Descriptive. 2 vols. 4to. Hf. mor.	10.00
POINSON. Eléments de Statique. 8vo. Hf. cf.	2.50
POISSON. Mécanique. 2 vols. 8vo.	6.00
PONCELET et LESBROS. Experiences Hydrauliques. 4to. Hf. mor.	4.00
PONTECOULANT. Système du Monde. 4 vols. 8vo. Hf. cf.	16.00
POUILLET. Eléments de Physique Experimentale. 3 vols. 8vo. Hf. cf.	7.50
SERRET. Cours d'Algèbre supérieure. 8vo. Hf. cf.	3.50
Eléments d'Arithmétique. 8vo. Hf. cf.	1.50
TERQUEM. Nouvelles Annales de Mathématiques. Monthly. Per annum	4.00
Nouvelles Annales de Mathématiques, 1856-7. 2 vols. Hf. cf.	6.50
Nouvelles Annales de Mathématiques. Tomes VIII. à XVII. Corresponding to the Years 1849-1859. 13 vols. Hf. cf.	
Bulletin Bibliographique, d'Histoire, et de Biographie Mathématiques. 8vo. Hf. cf.	3.00
TRESCA. Géométrie Descriptive. 8vo. Hf. mor.	3.00
VERHULST. Traité élémentaire des Fonctions Elliptiques. 8vo. Hf. cf.	3.50
VIELLE. Cours complémentaire d'Analyse et de Mécanique. 8vo. Hf. cf.	2.50

DODD'S MATHEMATICAL SERIES.

Text-Books for Colleges, Academies, and Schools.

BY PROF. JAMES B. DODD, TRANSYLVANIA UNIVERSITY, KY.

The attention of Instructors is respectfully invited to the books of Prof. Dodd's series, which are works of an established character, and used with satisfaction in prominent institutions of learning throughout the United States and Canada.

	RETAIL PRICE.
ELEMENTARY AND PRACTICAL ARITHMETIC. 12mo. 291 pp. Boards	\$ 0.50
HIGH-SCHOOL ARITHMETIC. 12mo. 364 pp. Cloth84
KEY TO ARITHMETIC. 12mo. 133 pp. Boards38
ELEMENTS OF ALGEBRA. 12mo. 217 pp. Sheep84
HIGH-SCHOOL ALGEBRA. (Table of Logarithms from 1 to 10,000. 16 pp.) 12mo. 306 pp. Sheep	1.25
KEY TO ALGEBRA. 12mo. 173 pp. Cloth84
ELEMENTS OF GEOMETRY AND MENSURATION. 12mo. 237 pp. Sheep	1.00
TRIGONOMETRY. With applications to SURVEYING, NAVIGATION, etc. 12mo. 212 pp. Tables 80 pp. Sheep	1.25

☞ A Primary Arithmetic, and books to complete the series, are in preparation, and will soon be published.

Dodd's Arithmetics and Geometry are adopted for the public schools of New York, and several other cities of the country.

Descriptions and many excellent commendations in our new Descriptive Catalogue of College and School Books, copies of which will be furnished gratis on application to

PRATT, OAKLEY, & CO., Publishers,

21 Murray Street, New York.

THE NEW AMERICAN CYCLOPÆDIA.

D. APPLETON & CO.,

Nos. 443 & 445 Broadway, New York,

HAVE JUST PUBLISHED

Vol. XI. of the

New American Cyclopædia,

A POPULAR DICTIONARY OF GENERAL KNOWLEDGE,

EDITED BY

GEORGE RIPLEY and CHARLES A. DANA,

Aided by a Numerous Select Corps of Writers from all parts of the United States.

This work is being published in about 16 large octavo volumes, each containing 750 two-column pages.

VOLS. I. TO XI. NOW READY.

To be followed by a new Volume every three months.

Price — In cloth, \$3; sheep, \$3.50; half morocco, \$4; half Russia, \$4.50 each.

From De Bow's Review, New Orleans.

"This great work, published by the Appletons of New York, has reached its eighth volume. It is one of the most valuable Cyclopædias in the world, and for Americans the most valuable. It is con-

tributed to by many of the ablest and best writers of the South, and is as fair to every section of the Union as it is possible for a work to be. It will be completed in about seven other volumes."

The following notice in *Silliman's American Journal of Sciences and Arts*, for March, is from the pen of the venerable senior editor, one of the highest living authorities on the subject:—

"In the American Cyclopædia the articles on Science are numerous and valuable, and elevate the work to the character of a compendium of modern science. These articles are in most cases written with decided ability, and evidently by persons who are familiar with the topics they discuss. While many of the less important subjects are presented with luminous brevity, others are more fully expanded. Among these are many topics of Natural History. Chemistry is presented with its equivalence and laws of combination, illustrated by many of its modern discoveries and practical applications; of the latter, an example is found in the full account of the manufacture of gelatine, of beer, and bricks, and in the ample history of gas-lighting. Neology, voltaic electricity, magnetism, and other departments of pure or applied science, are treated with reasonable fullness. As a literary work the Cyclopædia is written in a pure and chaste style, and exhibits the candor and fairness which should ever adorn a record of universal knowledge. L. S."

OUR SUBSCRIPTION LIST

Already numbers TWELVE THOUSAND names, from every section of the country, which evidences the universal popularity of this NATIONAL WORK.

THE NEW AMERICAN CYCLOPÆDIA

Is sold exclusively by subscription, and Agents have been appointed for almost all sections of the Union. In case, however, Agents are not to be found, we will receive subscriptions and forward copies of the work, free of expense, on receipt of the price. Persons at a distance may send money for one or more volumes at a time, it not being necessary to take the whole eight volumes at once, unless they feel inclined to do so. Those living in the city will find a subscription-book at our store, where their names will be entered, and the volumes, as published, sent to any part of the city.

GROVER & BAKER'S

Celebrated

SEWING MACHINES,

FOR

Family Use and Manufacturing Purposes.

OVER
40,000
IN DAILY USE
IN THE
United States



AND
Territories,
GIVING
UNIVERSAL
SATISFACTION.

THESE MACHINES sew from two spools; the thread requires no re-winding; and they make a STRONGER, HANDSOMER, AND MORE ELASTIC SEAM than any other Machine. They perform their work with a rapidity never before equalled, are almost noiseless in their operation, and are FAVORITES in every family where they have been introduced.

The "Scientific American," November 5th, 1859, says:—

"The machines manufactured by the Grover & Baker Sewing Machine Company are too well known by the public at large to need any recommendations at our hands, and we will simply add, that we have had one of them in use in our family for some time past, and it

is considered the most useful article in the house, next to the cradle, and no less indispensable than that."

"Zion's Herald," Nov. 16th, 1859, says:—

"We would recommend to families and to those about to organize a homestead, to be sure and have a Family Bible, and also one of Grover & Baker's SEWING MACHINES."

GROVER & BAKER SEWING MACHINE CO.,

18 Summer Street, Boston.
495 Broadway, New York.
730 Chestnut Street, Philadelphia.
181 Baltimore Street, Baltimore.

58 West Fourth Street, Cincinnati.
115 Lake Street, Chicago.
124 North Fourth Street, St. Louis.
91 Montgomery Street, San Francisco.

IMPORTANT EDUCATIONAL WORKS

RECENTLY PUBLISHED BY

GOULD & LINCOLN, BOSTON.

Lectures on Logic. By Sir William Hamilton, Bart., Professor of Logic and Metaphysics in the University of Edinburgh. With Notes from original materials, and an Appendix containing the latest development of his new Logical Theory. Edited by H. Longueville Mansel, B. D., Oxford, and John Veitch, A. M., Edinburgh. Royal 8vo. Cloth, \$3.00.

This volume completes the Metaphysical and Logical series of Sir William Hamilton, and will enable American scholars to possess themselves of the invaluable works of this eminent author, in two elegant octavo volumes, at six dollars,—half the price of the English edition.

Lectures on Metaphysics. By Sir William Hamilton, Bart. Edited by Prof. H. L. Mansel, B. D., Oxford, and John Veitch, M. A., Edinburgh. Royal 8vo. Cloth, \$3.00.

Hamilton's Lectures on Metaphysics have been received with eminent favor, and already adopted as a text-book in Harvard, Yale, Princeton, and other distinguished Colleges.

Prolegomena Logica. An Inquiry into the Psychological Character of Logical Processes. By H. Longueville Mansel, B. D., Professor of Moral and Metaphysical Philosophy, Oxford; Editor of Sir William Hamilton's Lectures; Author of "Limits of Religious Thought," etc. 12mo. Cloth, \$1.00.

This work is not an Introduction to Logic, but "an inquiry into the constitution and laws of the thinking faculty, such as they are assumed by the logician as the basis of his deductions." It is an important contribution to Mental Science, by one who stands in the front rank of living authors in this department of learning.

Limits of Religious Thought Examined. By H. Longueville Mansel, B. D., Professor of Moral and Metaphysical Philosophy, Oxford; Editor of Sir William Hamilton's Lectures, etc. 12mo. Cloth, \$1.00.

This able work of Prof. Mansel has passed through several editions, and is attracting marked attention among scholars and theologians, many of whom regard it as a defence of revealed religion unsurpassed since the appearance of Butler's Analogy.

Mental Philosophy; Including the Intellect, Sensibilities, and Will. By Joseph Haven, Professor of Intellectual and Moral Philosophy, Amherst College. Royal 12mo. Cloth, \$1.50.

Prof. Park, of Andover, says: "It is distinguished for its clearness of style, perspicuity of method, candor of spirit, acumen, and comprehensiveness of thought. I have been heartily interested in it."

"Earnestness, candor, and accuracy and elegance of diction characterize every page, and render the work as interesting to the general reader as it is valuable to the scholar."—D. H. COCHRAN, *Principal and Prof. of Moral and Intellectual Philosophy, State Normal School, Albany.*

This work has been extensively adopted as a text-book in Colleges and High Schools throughout the country.

Moral Philosophy; Including Theoretical and Practical Ethics. By Joseph Haven, D. D., late Professor of Moral and Intellectual Philosophy in Amherst College; Author of "Mental Philosophy." Royal 12mo. Cloth, \$1.25.

It is scientific in method and thorough in discussion, and its views on unsettled questions in morals are discriminating and sound. It treats largely of Political ethics,—a department of morals of great importance to American youth, but generally overlooked in text-books. In the history of ethical opinions, it is unusually rich and elaborate.

Elements of Moral Science. By Francis Wayland, D. D., late President of Brown University. 12mo. Cloth, \$1.25.

Elements of Political Economy. By Francis Wayland, D. D., 12mo. Cloth, \$1.25.

All the above works by Dr. Wayland are used as text-books in many of the colleges and higher schools throughout the Union, and are highly approved.

The Elements of Geology; Adapted to Schools and Colleges. With numerous Illustrations. By J. R. Loomis, President of Lewisburg University, Pa. 12mo. Cloth, 75 cents.

"It is surpassed by no work before the American public."—M. B. ANDERSON, LL. D., *President of Rochester University.*

Principles of Zoology; Touching the Structure, Development, Distribution, and Natural Arrangement of the RACES OF ANIMALS, Living and Extinct; with numerous Illustrations. For the use of Schools and Colleges. Part I. COMPARATIVE PHYSIOLOGY. By Louis Agassiz and Augustus A. Gould. Revised edition. \$1.00.

Thesaurus of English Words and Phrases; so classified and arranged as to facilitate the expression of ideas and assist in literary composition. By Peter Mark Roget, late Secretary of the Royal Society, London, etc. Revised and edited, with a List of Foreign Words defined in English, and other additions, by Barnas Sears, D. D., President of Brown University. A New American Edition, with Additions and Improvements. 12mo. Cloth, \$1.50.

The Earth and Man; Lectures on Comparative Physical Geography, in its relation to the History of Mankind. By Arnold Guyot. With Illustrations. 12mo. Cloth, \$1.25.

Prof. Louis Agassiz, of Harvard University, says: "It will not only render the study of Geography more attractive, but actually show it in its true light."

Essays in Biography and Criticism. By Peter Bayne, M. A., Author of "The Christian Life, Social and Individual." Arranged in two Series or Parts. 12mo. Cloth, \$1.25 each.

Guyot's Mural Maps. A series of Elegant Colored Maps for the Recitation Room, exhibiting the Physical Phenomena of the Globe. By Prof. Arnold Guyot, viz.: Map of the World, mounted, \$10.00. Map of North America, mounted, \$9.00. Map of South America, mounted, \$9.00. Map of Geographical Elements, mounted, \$9.00.

Chambers's Cyclopædia of English Literature. A Selection of the choicest productions of English Authors, from the earliest to the present time. Connected by a Critical and Biographical History. Two imperial octavo volumes, 700 pages each, double columns; with 300 elegant Illustrations. Embossed Cloth, \$5.00.

The Life of John Milton, Narrated in Connection with the Political, Ecclesiastical, and Literary History of his Time. By David Masson, M. A., Professor of English Literature, University College, London. Vol. I, 1608 to 1639. With Portraits, &c. Royal 8vo. Cloth, \$2.75.

An elaborate, erudite, and magnificent work.

Annual of Scientific Discovery for 1860; or, Year-Book of Facts in Science and Art, exhibiting the most important Discoveries and Improvements in Mechanics, Useful Arts, Natural Philosophy, Chemistry, Astronomy, Meteorology, Zoology, Botany, Mineralogy, Geology, Geography, Antiquities, &c. Edited by David A. Wells, A. M. 12mo. Cloth, \$1.25.

The Puritans; or the Court, Church, and Parliament of England, during the reigns of Edward VI. and Elizabeth. By Samuel Hopkins, Author of "Lessons at the Cross," etc. In 3 vols. Vols. I. and II. now ready. 8vo. Cloth, \$2.50 per vol.

This is a masterly work, by one of the most distinguished of living authors.

Cyclopædia of Anecdotes of Literature and the Fine Arts. Containing a copious and choice Selection of Anecdotes of the various forms of Literature, of the Arts, of Architecture, Engravings, Music, Poetry, Painting and Sculpture, and of the most celebrated Literary Characters and Artists, &c. By Kazlitt Arvine, A. M., Author of "Cyclopædia of Moral and Religious Anecdotes." With numerous Illustrations. 725 pp. 8vo. Cloth, \$3.00.

The Greyson Letters. Selections from the Correspondence of R. E. H. Greyson, Esq. Edited by Henry Rogers, Author of "The Eclipse of Faith." 12mo. Cloth, \$1.25.

"A book not for one hour, but for all hours; not for one mood, but for every mood; to think over, to dream over, to laugh over."—*Boston Journal.*
"The Letters are intellectual gems, radiant with beauty, happily intermingling the grave and the gay."—*Christian Observer.*

Knowledge is Power. A View of the Productive Forces of Modern Society, and the Results of Labor, Capital, and Skill. By Charles Knight. With numerous Illustrations. Revised, with Additions, by David A. Wells, Editor of the "Annual of Scientific Discovery." 12mo. Cloth, \$1.25.

[This is emphatically a book for the people. It contains an immense amount of important information, which everybody ought to be in possession of; and the volume should be placed in every family and in every School and Public Library in the land.]

Hugh Miller's Works. Seven volumes, uniform style, in an elegant box. Embossed Cloth, \$8.25.

Modern Text-Books in the Sciences.

A. S. BARNES AND BURR,
PUBLISHERS.

Chemistry, Geology, and Botany.

Porter's First Book of Chemistry and Allied Sciences.

Including an Outline of Agricultural Chemistry. By JOHN A. PORTER, A. M., Professor of Agriculture and Organic Chemistry at Yale College. 12mo. Cloth. Price, 50 cents.

Porter's Principles of Chemistry.

Embracing the most Recent Discoveries in the Science, and the Outlines of its Application to Agriculture and the Arts. Illustrated by numerous Experiments. By JOHN A. PORTER. 12mo. Cloth. Price, \$1.00.

This work has been prepared expressly for Public and Union Schools, Academies, and Seminaries, where an extensive course of study on this subject and expensive apparatus was not desired, or could not be afforded. A fair, practical knowledge of Chemistry is exceedingly desirable and almost a necessity at the present day, but it has been taught at very few Public or Union Schools, owing entirely to the want of suitable text-books, adapted to simple apparatus, or such as could be readily obtained. It is confidently believed that this work supplies this great want, and will be found in every respect just what is required.

Norton on Porter's First Book of Science.

Designed as an Introduction to the Study of Sciences. —PART I. Embraces Natural Philosophy and Astronomy, by WILLIAM A. NORTON, of Yale College. PART II. Chemistry, Physiology, Geology, and Agricultural Chemistry. Price, \$1.00.

Gregory's Hand-Book of Organic Chemistry.

For the use of Students. By WILLIAM GREGORY, Professor of Chemistry in the University of Edinburgh. Edited by J. M. SANDERS. 480 pp., 8vo. Cloth. Price, \$1.50.

Gregory's Hand-Book of Inorganic Chemistry.

For the use of Students. By WILLIAM GREGORY, Professor of Chemistry in the University of Edinburgh. Edited by J. M. SANDERS. 426 pp., 8vo. Cloth. Price, \$1.50.

These works are intended for the use of advanced students in this interesting science. They are the text-books in some of our leading colleges, and are well recommended by those who have used them.

Page's Elements of Geology.

By DAVID PAGE. Illustrated. 332 pp., 12mo. Cloth. Price, 75 cents.

The subject is here presented in its two aspects of interesting and important. Interesting, inasmuch as it exhibits the progressive conditions of the earth from the remotest periods, and reveals the character of the plants and animals which have successively adorned and peopled its surface; and important, as it determines the position of those metals and minerals upon which the arts and manufactures so intimately depend.

Emmons's Manual of Geology.

Designed for the use of Colleges and Academies. By EBENEZER EMMONS, State Geologist of North Carolina, late State Geologist of New York, Professor of Natural History and Geology in Williams College, etc. Illustrated with numerous engravings. 297 pp., 8vo. Cloth. Price, \$1.50.

A work that really requires nothing more than the name and reputation of the author to recommend it. Its intrinsic merit is such as must entitle it to a prominent position among kindred text-books.

Wood's Class-Book of Botany.

A new edition. Price, \$2.00.

Prof. Wood is now engaged in rewriting his celebrated Class-Book of Botany, embracing all the recent improvements, and will be unequalled by any now published. This splendid work will be ready January 1st, 1860.

Botany of the Southern States.

In Two Parts. PART I. Structural and Physiological Botany and Vegetable Products. PART II. Descriptions of Southern Plants. Arranged on the Natural System. Preceded by a Linnean and a Dichotomous Analysis. By JOHN DARBY, A. M., Professor of Natural Sciences in Auburn College, Ala. 612 pp., 12mo. Cloth. Price, \$1.00.

This work constitutes an outline of Botany in all its most important relations as a science. It has been the labor of the author for many years to bring it as near perfection as possible. It may be asserted that no work before the public affords so easy a means for the analysis of plants as this.

PUBLISHED BY A. S. BARNES & BURR,
51 and 53 John Street, New York.
Publishers of the National Series of Standard School-Books.

Modern Text-Books in the Sciences.

A. S. BARNES AND BURR,

PUBLISHERS.

Teachers and Scientific Men will please examine the following Text-Books, designed for Schools, Academies, and Colleges.

Single copies sent for examination AT ONE HALF THE RETAIL PRICE.

NATURAL PHILOSOPHY.

1. Norton's First Book of Natural Philosophy and Astronomy.

By WILLIAM A. NORTON, Professor of Civil Engineering in Yale College. 12mo. Cloth. Price, 50 cents.

The present little volume treats of the Elements of Natural Science, and is designed to meet the wants of young persons who do not intend to pursue a complete course of academical study. It is a good introduction to a more complete manual.

2. Ganot's Popular Physics.

By W. G. PECK. Introductory Course of Natural Philosophy, for the Use of Schools and Academies. Edited from "Ganot's Popular Physics." Price, \$ 1.00.

This work is illustrated by over 300 splendid engravings, and pronounced by all who have seen it as far superior to anything of the kind before the public. 10,000 copies have already been printed.

3. Peck's Elements of Mechanics.

Designed for Colleges, Academies, and High Schools. By WILLIAM G. PECK, M. A., Professor of Mathematics in Columbia College. 12mo. Cloth. Price, \$ 1.25.

A need has long been felt, by those who have classes in this science, of a text-book which may occupy an intermediate ground between those higher treatises which are found too difficult to be read with profit, save by a few, and too extensive to be studied, even by the few, in the limited time allotted to this branch of education, and those more elementary works, too simple for advanced classes, and, because of their non-mathematical character, not adapted to prepare the student for subsequent investigations in science. Professor Peck's work, just from the press, has been prepared with an especial view to meet this want. It will be found to contain a sufficient amount of information for those who lack either the leisure or desire to make mathematical sciences a specialty; and it will serve as a suitable introduction to those higher treatises on Mechanical Philosophy, which all must study who would appreciate and keep pace with the wonderful discoveries that are daily made in science.

4. Norton and Porter's First Book of Science.

Designed for Public and Private Schools. By W. A. NORTON and J. A. PORTER, Profes-

sors of Yale College. In two parts. Part I. embraces Natural Philosophy and Astronomy; Part II. Chemistry and Allied Sciences. 12mo. Cloth. Price of each part, separate, 50 cents; or two parts in one, \$ 1.00.

This work forms the most complete and interesting compendium of Familiar Science for young children now published. It embraces the principles of Natural Philosophy, Astronomy, Chemistry, Physiology, and Geology. The catechetical plan, found the most effective in impressing the minds of the younger class of pupils, has been adopted. Each subject is divested of the *dryness of detail* which is apt to characterize books of this grade, and is treated in a manner fitted to prepare the mind for future investigation, without wearying with present superfluity. It is thought to be eminently adapted to impart a practical knowledge of familiar things, and enable a child to understand the *why* and *wherefore* of the various facts and phenomena of every-day life.

5. A New Treatise on Astronomy and the Use of the Globes.

Containing Definitions, Motions, and Positions of the Sun, Moon, and Planets; Kepler's Laws, and the Theory of Gravitation; Gravitation; Refraction; Twilight and Parallax; Connections, Periods, Distances, Phenomena, and Magnitudes of the Heavenly Bodies composing the Solar System, &c.; also, an extensive collection of the most useful problems on the Globes, illustrated by a suitable variety of examples. By JAMES MCINTYRE, M. D., Professor of Mathematics and Astronomy in the Central High School of Baltimore. 12mo. Cloth. Price, 75 cents.

6. Natural Philosophy for Colleges.

By W. H. C. BARTLETT, LL. D., Professor of Natural Philosophy in the Military Academy of the United States.

BARTLETT'S SYNTHETIC MECHANICS	\$ 3.00
BARTLETT'S ACOUSTICS AND OPTICS	2.00
BARTLETT'S SPHERICAL ASTRONOMY	3.50
BARTLETT'S ANALYTICAL MECHANICS	3.50

7. Mrs. Willard's Astronomical Geography.

With the Use of the Globes. Arranged either for simultaneous Reading and Study in Classes, or for study in the common method. By EMMA WILLARD. Price, 75 cents.

PUBLISHED BY A. S. BARNES & BURR,

51 and 53 John Street, New York.

Publishers of the National Series of Standard School-Books.